



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>6</sup> : <b>C12Q 1/68</b></p>	<p><b>A2</b></p>	<p>(11) International Publication Number: <b>WO 99/63112</b> (43) International Publication Date: 9 December 1999 (09.12.99)</p>
<p>(21) International Application Number: PCT/US99/10940 (22) International Filing Date: 18 May 1999 (18.05.99) (30) Priority Data: 60/086,025 18 May 1998 (18.05.98) US Not furnished 17 May 1999 (17.05.99) US (71) Applicant (for all designated States except US): HUNT-WESSON, INC. [US/US]; 1645 West Valencia Drive, Fullerton, CA 92833 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROMICK, Thomas, L. [US/US]; 17637 Dandelion Lane, Chino Hills, CA 91709 (US). FRASER, Mark, S. [US/US]; 2331 E. Brookdale Place, Fullerton, CA 92831 (US). (74) Agents: CRAFT, Jeffrey, F. et al.; Pretty, Schroeder &amp; Poplawski, 19th floor, 444 South Flower Street, Los Angeles, CA 90071 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: PCR TECHNIQUES FOR DETECTING MICROBIAL AND VIRAL CONTAMINANTS IN FOODSTUFFS</p> <p>(57) Abstract</p> <p>A method of detecting the presence of living or dead microorganisms and viruses in a sample comprises adding to a predetermined volume of a sample comprising nucleic acid-containing microbe(s) and/or virus(es), known amounts of a pair of primers binding to sequences upstream and downstream to a universal or specific microbial and/or viral nucleic acid sequence and polymerase chain reaction (PCR) reagents, cycling the mixture to amplify the universal or specific microbial and/or viral nucleic acid sequence; adding a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PCR TECHNIQUES FOR DETECTING MICROBIAL AND VIRAL CONTAMINANTS IN FOODSTUFFS

### BACKGROUND OF THE INVENTION

5 This invention claims priority based on Serial No. 60/086,025, filed May 18, 1998.

#### Field of the Invention

The present invention relates to a novel technology provided in the form of novel primers, probes, and beacons, and their use in a rapid and accurate assay and kit for assaying for the presence of a microorganism or virus, particularly a bacterium, in a sample. More specifically, this invention relates to an accurate method for assessing microbial, e. g. bacterial or viral, contamination of sterile substances, such as sterile foods, by microorganisms and/or viruses relying on PCR and fluorescent beacon technologies.

#### Description of the Background

Beacons are fluorescent probes which contain short complementary sequences of nucleotide, (arms) attached to the 5' and 3' ends of a probe of a sequence complementary to a target nucleic acid. In addition, the beacon comprises a fluor, fluorophore or fluorogen (fluorogenic agent) and a quencher attached, e. g. via linkers, to the ends of the stem or arms. In the absence of a target, the fluor and the quencher remain close to one another held in place by a hairpin loop stem formed by hybridization of the arms. In this conformation, the beacon does not fluoresce because of quenching. However, when the segment encompassed by the loop hybridizes to a complementary sequence, the hybridization of the arms is prevented, the fluor, fluorophore or fluorogenic agent and the quencher are held apart and fluorescence appears. Thus, the appearance of fluorescence is an indication of hybridization of the probe (DNA internal segment of the beacon) to a complementary nucleic acid sequence of a target. The annealing of a strand of nucleic acid to its complement, thus, may now be measured by following changes in the physical properties of the nucleic acids which occur upon hybridization.

Prior technology utilized, and immobilized the hybridized strands onto, a solid surface, removed unhybridized probes and then determined the number of probes attached to the solid phase. The need for removal of unhybridized probes precluded the application

of solid phase hybridization to real time studies involving hybridization of nucleic acids. In addition, solid phase technology also has sensitivity limitations because the probes also bind non-specifically to the solid phase.

5 In the past the detection of microorganisms was made, generally, by phenotypic observation. Upon the realization that all living cells contain DNA and that DNA is responsible for the expression of phenotypic traits, novel detection methods relying on genetic parameters have been implemented. The polymerase chain reaction (PCR), a relatively recent technological development for amplification of DNA, provides significantly higher sensitivity and, thus, permits the detection of smaller quantities of nucleic acid by  
10 amplification and subsequent visualization, for instance, after electrophoretic separation and staining. Fluorescent dyes are also utilized by, for example, attachment to complementary oligomers (hybridization) that bind PCR-amplified DNA. These and other detection technologies eliminate the need for naked eye visualization of PCR products and, therefore, eliminate the need to electrophorese the nucleic acids, except for determining their molecular  
15 weight.

As is known in the art, the fidelity of specific DNA sequences produced by PCR is controlled by the primer DNA sequences used and by reaction conditions, such as thermocycling parameters, and the composition of the reaction mixture. DNA primers are generally designed to target specific sites vicinal to a desired sequence and applied to obtain  
20 DNA products of complementary sequences by the PCR, along with specifically tailored fluorescent molecular beacons designed to detect only the PCR product.

The detection of pathogens in edible products, cosmetics, medical fluids such as blood and IV solutions, and other products involved in commerce, is of great importance to avoid costly contamination, which may lead to outbreaks of disease and/or the need to discard large  
25 batches even after distribution. In most cases, even the etiological agent remains unidentified because of the sparcity of adequate testing technology to do this. Traditional technologies for recovering microorganisms from, for example, a foodstuff might include homogenization of the product in a buffered solution, inoculation of the homogenate in a selective enrichment medium, long hours of incubation at controlled temperatures, streaking of the broth onto a  
30 selective and/or differential agar medium, another incubation, isolation of colonies and finally a multiplicity of tests for biochemical or immunological characteristics and microscopic examination. Virulence is mostly tested on pure cultures which require several

days of incubation. In addition, for sterility testing, the mere detection of biologically active DNA is critical.

Accordingly, there is a need for a rapid, simple, inexpensive and sensitive method for the general detection of microorganisms and/or viruses in commercial products as well as in pure cultures of microorganisms and viruses.

### SUMMARY OF THE INVENTION

This invention relates to an in vitro method of detecting the presence of a microbe or a virus in a food sample, comprising the steps of:

(a) forming a polymerase chain reaction mixture by combining (1) a predetermined volume of a food sample to be tested for the presence of a nucleic acid sequence comprising a universal or specific nucleic acid sequence indicative of a microbe or a virus and sequences upstream and downstream of the universal or specific nucleic acid sequence, (2) known amounts of a first nucleic acid primer and a second nucleic acid primer for binding to the upstream sequence and the downstream sequence, respectively, and (3) polymerase chain reaction reagents;

(b) forming a polymerase chain reaction product by cycling the polymerase chain reaction mixture under conditions effective to amplify the universal or specific nucleic acid sequence, if present, to replicate and attain about 0.25 to about 10,000 µg nucleotide product/µl mixture;. and

(c) determining whether or not the universal or specific nucleic acid sequence is present in the polymerase chain reaction product, the presence of the universal or specific nucleic acid sequence being indicative of the presence of a microbe or a virus in the food sample.

Any suitable method is used to determine the presence of the universal or specific nucleic acid sequence is present in the polymerase chain reaction product. Fluorescent intercalating reagents, such as ethidium bromide, can be used in conjunction with a fluorescence detection system. Alternatively, suitable oligonucleotide primers(s) that are hybridizably complementary to at least a portion of the reaction product, and bearing a fluorescent label and/or quencher can also be used in conjunction with the PCR amplification and/or fluorescence detection system as known in the art.

This invention also relates to an in vitro method of detecting the presence of a microbe

or a virus in a food sample, comprising:

adding to a pre-determined volume of a sample suspected of comprising nucleic acid-containing microbe(s), known amounts of a pair of primers binding to sequences up-stream and down-stream to a universal or specific microbial and/or viral nucleic acid sequence and  
5 polymerase chain reaction (PCR) reagents to form a mixture;

cycling the mixture about under conditions effective to amplify the universal or specific microbial and/or viral nucleic acid sequence to attain about 0.25 to about 10,000  $\mu\text{g}$  universal microbial nucleic acid/ $\mu\text{l}$  mixture;

adding a polynucleotide comprising a DNA internal segment that is hybridizably  
10 complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another. Optionally, the terminus of one arm segment is operatively connected to a  
15 fluorogen and the terminus of the other arm segment is operatively connected to a quencher, such that the polynucleotide comprises a "fluorescent beacon." The polynucleotide can form a stem-loop structure when there is no universal or specific microbial and/or viral nucleic acid present in the sample. A change in conformation of the stem-loop structure is detected by a fluorescent detection system; and detecting fluorescence emitted by the sample. Any  
20 suitable reagent containing a fluorogen or fluorogenic agent, such as fluorescein, or an intercalating agent, such as ethidium bromide, can be used in detecting a conformational change in the polynucleotide. The reagent can also be a suitable oligonucleotide bearing a fluorescent label.

Alternatively, the polynucleotide is a "fluorescent beacon." When no universal or  
25 specific microbial and/or viral nucleic acid is present and replicated by PCR in the sample, the DNA internal segment of the fluorescent beacon is single stranded, the complementary sequences of the arms are hybridized to one another and the fluorogenic agent is quenched by the quencher; when there is microbial and/or viral DNA in the sample, a portion of the universal or specific microbial and/or viral nucleic acid is hybridized to the internal segment  
30 of the beacon, and the fluorogenic agent fluoresces.

This invention also relates to a kit for detecting a microbe or virus. The detection kit comprises primers binding to sequences up-stream and down-stream to universal or specific

microbial and/or viral nucleic acid sequence (s) and polymerase chain reaction (PCR) reagents; a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another. Optionally, the terminus of one arm segment is operatively connected to a fluorogen and the terminus of the other arm segment is operatively connected to a quencher, such that the polynucleotide comprises a "fluorescent beacon." The polynucleotide can form a stem-loop structure when there is no universal or specific microbial and/or viral nucleic acid present in the sample. A change in conformation of the stem-loop structure is detected by a fluorescent detection system. The kit also contains instructions for use of the kit to detect the presence of a microbe (s) and/or virus (es), in particular bacteria, and the kit optionally contains PCR reagents.

#### 15                    **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

This invention arose from the desire by the inventors to improve on prior art technology for assessing in a short time whether a sample is contaminated with microorganisms or viruses in an accurate and simple manner. This invention provides an assay and detection kit combining the benefits of nucleic acid amplification by PCR, the specificity of DNA hybridization probes and the signal amplification capability of fluorescent molecular beacon technologies to detect the presence of microorganisms and/or viruses in a product without needing to separate unbound beacon molecules.

This invention relates to an in vitro method of detecting the presence of a microbe or a virus in a food sample, comprising the steps of:

- 25            (a)    forming a polymerase chain reaction mixture by combining (1) a predetermined volume of a food sample to be tested for the presence of a nucleic acid sequence comprising a universal or specific nucleic acid sequence indicative of a microbe or a virus and sequences upstream and downstream of the universal or specific nucleic acid sequence, (2) known amounts of a first nucleic acid primer and a second nucleic acid primer for binding to the upstream sequence and the downstream sequence, respectively, and (3) polymerase chain  
30            reaction reagents;

(b) forming a polymerase chain reaction product by cycling the polymerase chain reaction mixture under conditions effective to amplify the universal or specific nucleic sequence, if present, to replicate and attain about 0.25 to about 10,000 µg nucleotide product/µl mixture;. and

5 (c) determining whether or not the universal or specific nucleic acid sequence is present in the polymerase chain reaction product, the presence of the universal or specific nucleic acid sequence being indicative of the presence of a microbe or a virus in the food sample.

Any suitable method is used to determine the presence of the universal or specific  
10 nucleic acid sequence is present in the polymerase chain reaction product. Fluorescent intercalating agents, such as ethidium bromide, can be used in conjunction with a fluorescence detection system. Alternatively, a suitable oligonucleotide primer(s) that is hybridizably complementary to at least a portion of the reaction product, and bears a fluorescent label and/or quencher can also be used in conjunction with the PCR amplification  
15 or the fluorescence detection system. For example, when the universal or specific nucleic acid sequence is 5'-TAGAAGC-3', suitable oligonucleotide primers(s) for purposes of amplification and/or detection of amplification reaction products include 5'-GCTAAGGTCCCAAAGT-3' , which is usefully labeled with a fluorogen, such as fluorescein, at its 5'-end, and/or 5'-ACTTTGGGACCTTAGC-3' , which can also be usefully  
20 labeled at its 3'-end with a quencher, such as dabcy1, to quench unhybridized fluorogen-labeled primer in the reaction mix.

The present method also provides a broadly applicable in vitro method of detecting the presence of one or more microbe (s) and/or virus (es) in a sample, which comprises

25 adding to a pre-determined volume of a sample suspected of comprising nucleic acid-containing microbe(s) known amounts of a pair of primers binding to sequences up-stream and down-stream to a universal microbial and viral nucleic acid sequence and polymerase chain reaction (PCR) reagents to form a mixture;

cycling the mixture under conditions effective to amplify the universal microbial and/or viral nucleic acid sequence to attain about 0.25 to about 10,000 µg universal microbial  
30 nucleic acid/µl mixture;

adding a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a



first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another. Optionally, the terminus of one arm segment is operatively connected to a fluorogen and the terminus of the other arm segment is operatively connected to a quencher, such that the polynucleotide comprises a "fluorescent beacon." The polynucleotide can form a stem-loop structure when there is no universal or specific microbial and/or viral nucleic acid present in the sample. A change in conformation of the stem-loop structure is detected by a fluorescent detection system and detecting any fluorescence emitted by the sample, wherein when no universal microbial and/or viral nucleic acid is present and replicated by PCR in the sample the internal DNA segment remains single stranded. Any suitable reagent containing a fluorogen or fluorogenic agent, such as fluorescein, or ethidium bromide can be used in detecting a conformational change in the polynucleotide. The reagent can also be a suitable oligonucleotide primer, bearing a fluorescent label or quencher, as described above.

If the polynucleotide is a "fluorescent beacon," the complementary sequences of the extension arms are hybridized to one another and the fluorophore or fluorogenic agent is quenched by the quencher, and when there is microbial and/or viral DNA in the sample, its universal microbial and/or viral nucleic acid sequence is hybridized to the internal segment of the beacon and the fluorogenic agent fluoresces.

As already described above, the polynucleotide (or beacon) is a single stranded nucleic acid molecule possessing a stem and loop structure, where the loop is a probe with a sequence which is complementary to a pre-determined sequence in a target nucleic acid. The stem, on the other hand, is formed by the annealing of two complementary arm segment sequences placed on both sides of the probe segment (DNA internal segment of the polynucleotide) and which DNA arm segment sequences are unrelated to the target sequence to avoid hybridization. Typically, the 5' and 3' ends of the arms are attached, respectively, to a fluorescent moiety and to a non-fluorescent quenching moiety to form a "fluorescent beacon" structure. Thus, the stem portion of the stem and loop structure keeps these two moieties in close proximity and, thereby, quenches any fluorescence by the fluor, fluorophore or fluorogenic agent. When the probe segment encounters a target molecule and hybridizes to it to form a more stable, double stranded structure than the stem and loop structure. The existence of the two structures is precluded by the relative rigidity of nucleic acid structures.

Thus, when the probe encounters a complementary sequence, it undergoes a spontaneous conformational change that opens up the stem and forces the arms apart from one another, and, thereby, causes the separation of the fluorophore or fluorogen (or fluorogenic agent) and the quencher and permits, upon incidence of ultraviolet light, the fluorophore to fluoresce.

5 Moreover, because unhybridized beacons are quenched, it is not necessary to remove them from the medium to detect by fluorescence the presence of specific nucleic acids in homogeneous phase and in living cells.

In one preferred embodiment, the DNA internal segment and the two DNA arm segments are complementary to a contiguous universal microbial nucleic acid sequence, and  
10 more preferably the DNA internal segment is about 7 to 19 nucleotides long, more preferably about 9 to 15, and still more preferably about 11 to 13 nucleotides long, each DNA arm segment is preferably about 5 to 9 nucleotides long, more preferably about 6 to 8 nucleotides long, and still more preferably 7 nucleotides long, and each primer is preferably about 10 to 20 nucleotides long, more preferably about 12 to 18, and still more preferably about 14 to  
15 16 nucleotides long. In one of the most preferred embodiments the polynucleotide comprises one arm segment with at least one C or G nucleotide(s) at the 5' terminus and the other arm segment with a similar number of complementary nucleotide(s) of the 3' terminus. In this manner, the arm segments strongly hybridize by formation of at least one G:C pair to, thereby, prevent sliding of the DNA segment. The DNA arm segments comprise any  
20 possible combination of C, A, T and G nucleotides, and preferably about 1 to 8 C or G at the 5' terminus and a similar number of complementary nucleotides at the 3' terminus, more preferably about 1 to 3 C or G nucleotide at the 5' terminus and 1 to 3 complementary nucleotides at the 3' terminus, still more preferably 2 to 6 C or G nucleotides at the 5' terminus and 2 to 6 complementary nucleotide at the 3' terminus. In addition, the  
25 polynucleotide's arm segments may further comprise at least one A or T nucleotide(s) vicinal to the C or G nucleotide(s) located at the 5' terminus in a similar number of complementary nucleotide(s) vicinal to the G or C nucleotide(s) at the 3' terminus. In this manner, the hybridization of the arm segments is further reinforced by formation of one or more A:T pairs next to the one or more terminal G:C pairs, preferably the polynucleotide's arm  
30 segments comprise 1 to 4 A or T nucleotide vicinal to the C or G nucleotide at the 5' terminus and 1 to 4 complementary nucleotide(s) thereto which are vicinal to the G or C nucleotide at the 3' terminus, and still more preferably about 2 to 3 A or T nucleotides and

2 to 3 complementary nucleotides.

In another preferred embodiment of the invention, the DNA arm segments are complementary to one another, but at least one of them is not hybridizably complementary the universal microbial or viral nucleic acid sequence. Such non-complementary arm segments are known as "false arms." In this embodiment, the DNA internal segment is preferably about 7 to 19 nucleotides long, more preferably 10 to 15 nucleotides long, and still more preferably 12 to 14 nucleotides long, each DNA arm segment is preferably 5 to 9 nucleotides long, and more preferably 6 to 8 nucleotides long, and each primer is about 10 to 20 nucleotides long, and more preferably 12 to 18 nucleotides long. However, in some instances, longer or shorter sequences are also suitable. In this case, the polynucleotide's DNA arm segments may comprise one or more C or G nucleotide(s) at the 5' terminus and a similar number of complementary nucleotide at the 3' terminus. The pairing of these nucleotide(s) provides strong hybridization and prevents the sliding of the DNA segments with respect to one another. A preferred number of C or G nucleotides at the 5' terminus of the polynucleotide's (or fluorescent beacon's) arm segments is about 1 to 8, and more preferably about 3 to 5 C or G nucleotides, and a similar number of complementary nucleotide at the 3' terminus of the extension arms. In addition, the arm segments can further comprise one or more A or T nucleotide vicinal to the one or more C or G nucleotide(s) at the 5' terminus in a similar number of complementary nucleotides vicinal or next to the one or more G or C nucleotide(s) at the 3' terminus. As may be surmised, the addition of A:T pairs next to the G:C pairs further reinforces the strength of the hybridization exhibited by the extension arms. The number of A or T nucleotides vicinal to the C or G nucleotides at the 5' terminus of the extension arms is preferably 1 to 4, and more preferably 2 to 3. Consequently, the number of complementary nucleotides next to the G or C nucleotides at the 3' terminus is also 1 to 4, and more preferably 2 to 3.

The reagents, including enzymes, which may be utilized in the polymerized chain reaction (PCR) are known in the art. See, for example, US Pat. Nos. 5,210,015; 4,683,195; 4,683,202; 4,965,188; 4,800,159; and 4,889,818, the relevant portions of which are incorporated by reference. However, preferred cycling conditions for this reaction are about 1 to 30 cycles, more preferably 5 to 20 cycles, and still more preferably 10 to 15 cycles, at alternating between temperatures of about (1) 58 to about 95°C: to (2) about 58 to about 95°C. Some of the more preferred temperature combinations being about 95: about 58°C,

about 58: about 74°C and about 74: about 95°C. However, other combinations are also suitable. A preferred amount of beacon DNA is approximately the same as or about the concentration of both primer DNAs used. However, other proportions may also be utilized as an artisan would know. See, for example, the US Patents listed above.

5 In another preferred embodiment, the DNA internal segment of the polynucleotide binds to universal or conserved nucleic acid sequences of a large number of viruses and microorganisms, including bacteria, yeast, molds and protista. In still a more preferred embodiment, the internal DNA segment comprises a sequence which hybridizes to, and may be complementary to a contiguous specific universal bacterial nucleic acid sequence and, therefore, it hybridizes to all classes of bacteria. By means of example, a bacteria-specific nucleic acid sequence comprises 5'-ATCTTCG-3' (SEQ. ID NO: 115), and the internal DNA segment comprises a complementary sequence thereto, i.e. 3'-TAGAAGC-5' (SEQ. ID NO: 116). Other sequences, however, may also be utilized. In one preferred embodiment, both the internal DNA segment and the extension arms are formed by a sequence which is complementary to a contiguous bacteria-specific bacterial nucleic acid sequence. By means of example, the nucleic acid sequence of the polynucleotide (or beacon) may comprise 3'-GGTGGCTTAGAAGCAGCCACC-5' (SEQ. ID NO: 117). Others, however, are also suitable.

15 In yet another embodiment, the polynucleotide (or fluorescent beacon) may be a specific probe for a class of microorganisms, for a specific species or for a group of species, depending on the DNA internal segment chosen. Examples of all of the above are provided in the Tables below.

20 The present method is similarly applicable to the detection of viruses by implementing the method described above. Similarly, nucleic acid sequences which are specific for yeast may be utilized to design primers which will bind upstream and downstream from the sequences to amplify the yeast nucleic acid sequence, and to manufacture a polynucleotide (or fluorescent beacon) by selecting a complementary DNA sequence which includes the DNA internal segment and the DNA arm segments and may additionally have the A:T and C:G termini. Alternatively, the DNA internal segment may comprise a sequence complementary to the yeast nucleic acid and the arm segments may be "false arms" constituted by complementary strings of A, T, C and G for form A:T and C:G pairs. This may also be attained with specific sequences for molds and/or protista in the same manner.

Various universal and specific microbial and viral nucleic acid sequences, primers, and DNA sequences suitable for use in the polynucleotide (or fluorescent beacon) in accordance with the present invention are provided below. However, the practice of this invention is not limited to the sequences described herein but, in addition, may be practiced with other universal nucleic acid sequences or sequences specific for either a species or a class of microorganism or virus. In another embodiment of the invention, whether or not a sample is sterile or has microbial and/or viral contamination may be determined by using a "pre-mix" of reagents with which a sample is mixed, and a specialized instrument for detecting fluorescence. This embodiment may be performed in a period of time as short as about 10 minutes. In this embodiment, the reagent "pre-mix" may comprise two primers and a fluorescent molecular probe or beacon, to which are added the sample and the PCR reagents, e.g. enzyme.

The necessary elements for implementing this technology are provided as a kit for detecting a microbe and/or virus. The detection kit comprises in separate containers primers binding to sequences up-stream and down-stream to a universal or specific microbial and/or viral nucleic acid sequence; and optionally, a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another. Optionally, the terminus of one arm segment is operatively connected to a fluorogen and the terminus of the other arm segment is operatively connected to a quencher, such that the polynucleotide comprises a "fluorescent beacon."

Also included in the kit are instructions for use of the kit to detect the presence of a microbe(s) or virus(es).

As described above, in one preferred embodiment the DNA internal segment and the two arm segments of the polynucleotide (or fluorescent beacon) are fully complementary to a contiguous universal microbial nucleic acid sequence. Preferably, the internal DNA segment is about 7 to about 19 nucleotide long, each extension arm is about 5 to about 9 nucleotides long, and each primer is about 10 to about 20 nucleotides long. In another embodiment, only

the DNA internal (or probe) segment of the polynucleotide (or fluorescent beacon) is complementary to a universal microbial nucleic acid sequence. Although the DNA arm segments are complementary to one another, one or both of them can be unrelated to the universal microbial nucleic acid sequence. In the latter case, although the length of the different parts of the fluorogenic beacon are similar to the one described above, the arm segments preferably comprise at least one C or G nucleotide at the 5' terminus in a similar number of complementary nucleotide at the 3' terminus, which permits the formation of G:C pairs by hybridization. More preferably the arm segments comprise about 1 to 8 C or G nucleotide at the 5' terminus and 1 to 8 complementary nucleotide(s) at the 3' terminus. Still more preferably 2 to 6 G or C nucleotide(s) at the 5' nucleotide at the 5' terminus and 2 to 6 complementary nucleotide at the 3' terminus. In addition, the DNA arm segments may further comprise one or more A or T nucleotide(s) vicinal or next to the C or G nucleotide at the 5' terminus, and the other arm comprises a similar number of complementary nucleotides vicinal to the G or C nucleotide(s) at the 3' terminus, thereby reinforcing the strength of the double stranded DNA portion. Preferably, the A or T nucleotide segments comprises 1 to 4 nucleotide(s) vicinal to the C or G nucleotide(s) at the 5' terminus and the other arm segment comprises 1 to 4 complementary nucleotide(s) to the A or T, which are placed vicinal or next to the G or C nucleotide(s) at the 3' terminus. In another embodiment, the DNA internal segment is complementary to a bacteria-specific nucleic acid sequence. Although the DNA arm segments are still complementary to one another, one or both can be unrelated to the bacteria-specific nucleic acid sequence. In this case, one of the DNA arm segments of the polynucleotide (or beacon) comprises at least one and up to 8 C or G nucleotide(s) at the 5' terminus and the other arm segment comprises a similar number of complementary nucleotides at the 3' terminus. As already explained above, the DNA arm segments can additionally comprise 1 to 4 A or T nucleotides(s) next to the C or G nucleotide(s) at the 5' terminus and the other arm comprises 1 to 4 complementary nucleotide vicinal to the one or more G or C nucleotide(s) at the 3' terminus. Examples of sequences suitable for use as arm segments are 5'- GCTAG -3' (SEQ. ID NO: 118) and 3'- CGATC -5' (SEQ. ID NO: 119), although many others are also suitable. By means of example, the internal DNA segment of the fluorogenic beacon may be complementary to a conserved region of a small ribosomal RNA (rRNA) operon, a first primer may comprise 5'-GCTAAGGTCCCAAAGT-3' (SEQ. ID NO: 120), and a second primer may comprise 5'-AGAACGCTCTCCTACC-3' (SEQ. ID NO: 121). Another

example is that where the DNA internal segment of the polynucleotide (or fluorescent beacon) is complementary to a conserved region of the small rRNA operon encoding a 23S rRNA, wherein the internal DNA segment of the beacon is complementary to the sequence 5'- 5'-CTTAGAAGCAG -3' -3' (SEQ. ID NO: 122). The oligonucleotide primers of the present invention, useful in the present kit, are optionally connected at their 5'-ends to a fluorogen, e.g., fluorescein, or at their 3'-ends to a quencher, e.g., dabcyI.

The universal or specific nucleic acid sequence may be selected, and the complementary internal DNA segment of the polynucleotide (or fluorescent beacon) designed, for microorganisms such as bacteria, yeast, mold or protista. The DNA internal segment may be complementary to a contiguous nucleic acid sequence which is universal to a class of microbes as opposed to all microbes. Examples of these are provided below. The DNA internal segment moreover, may be complementary to a contiguous nucleic acid sequence which is virus-specific or specific to a class of viruses. When the kit is applied to the detection of bacteria at large, the DNA internal segment can comprise a sequence that is complementary to a contiguous universal bacteria-specific bacterial nucleic acid sequence, such as 5'-TAGAAC-3', or its complement, although others are also suitable. In another embodiment both the DNA internal segment and the arm segments comprise a nucleic acid sequence, such as 3'-GGTGGCTTAGAAGCAGCCACC-5', although others may also be utilized. Where the present technology is applied to the detection of virus, the DNA internal segment or both the DNA internal segment and the arm segments may be complementary to a nucleic acid which is conserved in all viruses, conserved over a class of viruses or a virus-specific nucleic acid sequence.

Any fluorogen (or fluorogenic agent) known in the art may be utilized with the present technology. Particularly suitable are pico green, fluorescein, edans, and the like. However, others may also be utilized. Similarly, any quencher known in the art is suitable for use herein. By means of example, dabcyI is suitable for the present purpose, although others may be utilized. Both the fluorogenic agent and the quencher are linked to the DNA portion of the fluorogenic beacon by C<sub>6</sub>-thio or C<sub>7</sub>-amino linkers known in the art, and the linkage is conducted as is known in the art. See, for example, Columns 1-2, page 1143, Schofield et al., Appl. & Ennison. Microbiol. 63(3):1143 (1997); Tyagi & Kramer, Nature Biotech. 14:303 (1996), the text of which is incorporated herein by reference.

In one preferred embodiment of the present kit, the primers and the polynucleotide

(or fluorescent beacon) are pre-mixed in bulk, or in unitary amounts and provided in unitary form. The kit may also comprise one or more controls such as universal microbial and viral nucleic acid sequences and/or microbe-specific and viral-specific nucleic acid sequences. The controls may be provided as a panel for all microorganisms and/or viruses as well as  
5 different classes or specific species of bacteria or viruses.

In a preferred embodiment of this invention, the DNA primers comprise universal DNA oligomers, which are suitable for amplifying all eukaryotic and prokaryotic microorganisms, including bacteria, mold, yeast and/or protista, or viruses. In a broadly encompassing embodiment of the method of this invention, a predetermined volume of  
10 sample is mixed with the primers and other PCR reagents, a PCR reaction conducted, a fluorogenic beacon added and fluorescence detected.

The kit may further comprise PCR reagents. A most preferred embodiment of the kit, comprises a "pre-mix" of reagents, and instructions for DNA amplification and for conducting a PCR under specific cycling conditions. As already indicated, the kit of the  
15 invention may be a kit for generally assessing the presence or absence of microorganisms, e.g., bacteria, in a sample or may be custom tailored for the detection of specific organisms, such as any desired bacterium. An embodiment of the method and kit of the invention suitable for the determination of bacterial contamination in general may comprise a bacteria-specific nucleic acid sequence and its corresponding primers. The inventor has found that  
20 when a bacteria-specific sequence and its primers are subjected to PCR under certain conditions, the method of the invention provides high specificity for the bacterial genus. The universal microbial and/or viral nucleic acid sequence is preferably provided in the form of a stem-loop structure. However, the nucleic acid sequence need not be in this form. The nucleic acid target sequence may be in linear and/or circular form, as well. The  
25 polynucleotide (or fluorescent beacon), however, must generally conform to a stem-loop structure, and must comprise at least an DNA internal segment which is complementary to the target sequence. See, Tyagi and Kramer, Nature Biotech. 14:303 (1996). If the microbial and/or viral target sequence is not in the form of a stem-loop structure, the polynucleotide (or fluorescent beacon) must contain additional nucleotides at either, or both,  
30 end(s) of the complementary region to form a "false stem-loop" (arm segments). See, Schofield et al., App. & Environmental Microbiol. 63(3):1143 (1997).

A preferred "false stem-loop" comprises one or more of G:C, C:G, T:A, A:T, G:C,



b/e last and/or C:G 1st, and the false stem-loop sequence is generally about 3 to about 51 nucleotides long, and preferably about 7 to about 31, and more preferably about 12 to about 28 nucleotides long. Most preferred are false stem-loop segments about 13, about 15, about 17, about 19 and about 21 nucleotides long. However, other lengths are also suitable. The nucleotide segments may also comprise even numbers of nucleotide. However, odd-number nucleotide segments are preferred because they provide a trans-hybridization configuration which yields greater fluorescence. Where a preferred false stem structure may not be attainable, the nucleotide segment of the polynucleotide (or beacon) must comprise at least about 2 G:C nucleotide pairs positioned at the termini (linker arm ends), and up-stream from it, preferably immediately up-stream, at least one T:A nucleotide pair(s) to prevent slipping in the stem hybrid (G:C, C:G, T:A,...) for proper attachment of the polynucleotide's (or beacon's) arms where in single stranded form. This sequence is generally followed by at least about 1, preferably about 2 more complementary base pairs, and up to about 8 complementary base pairs, preferably G:C pairs, to require a high enough melting temperature for the stem hybrid structure to remain stable at ambient temperature.

A preferred polynucleotide (or beacon molecule) for the universal detection of bacteria comprises 5'-GGTGGCTGCTTCTAAGCCACC-3' or analogues thereof which hybridize under the following conditions, and preferably under stringent conditions, to a DNA sequence complementary thereto. Other polynucleotide DNAs (or beacon molecules) are also suitable and may be utilized in conjunction with this invention. Such molecules are designed to be complementary to conserved regions in the small rRNA operon encoding the 23S rRNA. Its corresponding primers comprise segments 5'-GCTAAGGTCCCAAAGT-3', and 5'-AGAACGCTCTCCTACC-3'. These sequences were deduced from highly conserved regions of the rRNA operon encoding 23s rRNA. Other universal probes or internal DNA segments suitable for use with this invention are those which have a sequence complementary to the target nucleic acid sequences described by Hill (1996), supra, and include highly conserved sequences such as a 16S rRNA, an rRNA spacer, a 16S rDNA, a 16S DNA, which are listed in Table 2 of the publication. Hill (1996), supra also provides in Table 2 sequences specific for individual microorganism targets, including *Aeromonas hydrophila* aer, *Brucella* spp. 16S rDNA, *Brucella* spp. Putative OMP, *Campylobacter jejuni* flaA, *E. coli* flaA, *C. jejuni* 16S rRNA, *E. coli* 16S rRNA, *C. Lari* 16S rRNA, *Clostridia* 16S rDNA (2 sequences), *Clostridium botulinum* Neurotoxin A, *Clostridium botulinum* Neurotoxin B,

etc. Some of the sequences encompassed by the group to which the invention is applicable are listed below. This, however, is not an all inclusive list of known highly conserved and specific sequences for microbial and viral targets.

**Table 1: Examples of Highly Conserved Target Sequences**

Organism	Target (bp)	Size	Primer Sequences	Ann.Temp. (°C)	Cycles
Bacteria	16S rRNA	408	CAGCMGCCGCGGTAATWC CCGTCAATTCMTTTRAGTTT	50	30
"	rRNA Spacer	Varies	GAAGTCGTAACAAGG CAAGGCATCCACCGT	55	25
"	16S rDNA	-335	GAGTTTGATCCTGGCUCA	60	35

Table 2 below shows several sequences of molecular beacons and indicates whether the sequences are universal or specific for a certain organism. See, Hill, W. E., Crit. Rev. Food Sci. & Nutrition 36(1 & 2):123 (1996).

**Table 2: Examples of Molecular Beacon Sequences**

Organism	Molecular Beacon Sequence
Universal	5'-CCTGCACGGGCGGTGTGTACGCAGG-3' (SEQ. ID NO: 123)
R. Albus 8	5'-CCCCCGTCATGCGGCTTCGTTATGGGGG-3' (SEQ. ID NO: 124)
F.succinogS85	5'-GCTGCCTGCCCTGAACTATCCAAGAGGCAGC-3' (SEQ. ID NO: 125)

Examples of sequences for organism specific targets and the respective organism are provided in Table 3 below.

**Table 3: Examples of Specific Target Sequences**

Organism	Target (bp)	Size	Primer Sequences	Ann.Temp. (°C)	Cycles
All bacteria	16S rRNA	408	CAGCMGCCGCGGTAATWC CCGTCAATTCMTTTRAGTTT (SEQ ID NO:1)	50	30

**Table 3: Examples of Specific Target Sequences (Cont'd)**

5	All bacteria	rRNA spacer	Variable	GAAGTCGTAACAAGG CAAGGCATCCACCGT (SEQ ID NO:2)	55	25
	All bacteria	16S rDNA	-335	GAGTTTGATCCTGGCUCA CTGCTGCCTCCCGTA (SEQ ID NO:3)	60	35
	Eubacteria	16S DNA	-1500	AGAGTTTGATCCTGGCTCAG AAGGAGGTGATCCAGCCGA (SEQ ID NO:4)	55 42	26 25-30
	<i>Aeromonas hydrophila</i>	<i>aer</i>	209	CCAAGGGGTCTGTGGCGACA TTTCACCGGTAACAGGATTG (SEQ ID NO:5)	55	30
	<i>Brucella</i> spp.	16S rDNA	801	TGCTAATACCGTATGTGCTT TAACCGCGACCGGATGTCAA (SEQ ID NO:6)	50	?
10	<i>Brucella</i> spp.	Putative OMP	635	Sequences not reported	60	50
	<i>Campylobacter jejuni</i> and <i>C. coli</i>	<i>flaA</i>	560	ATGGGATTTTCGTATTAAC GAACTTGAACCGATTG (SEQ ID NO:7)	37	25
	<i>C. jejuni</i> , <i>C. coli</i> , <i>C. lari</i>	16S rRNA	426	ACCTTGTTACGACTTCACCCCA GAGAGTTTGATCCTGGCTCAG (SEQ ID NO:8)	52	40
15	<i>Clostridia</i>	16S rRNA	-1371	GATCCTGGCTCAG GACGGGCGGTGTGTACAA (SEQ ID NO:9)	50	40
	<i>Clostridia</i>	16S rDNA	-502	AAACTCAAATGAATTGACGG GACGGGCGGTGTGTACAA (SEQ ID NO:10)	50	40
20	<i>Clostridium botulinum</i>	Neurotoxin B	-1500	GATGGAACCACCATTTGCWAG WACATCWATACAWATTCCTGG (SEQ ID NO:11)	37	25
	<i>Clostridium botulinum</i>	Neurotoxin A	-1340	GATCCTGTAAATGGTGTTG CAAGTCCCAATTATTATAACTTTG AT (SEQ ID NO:12)	55	35
	<i>Coliforms</i>	<i>lacZ</i>	?	TGAAAGCTGGCTACAGGAAGGCC GGTTTATGCAGCAACGGACGTCA (SEQ ID NO:13)	60	30
	<i>Entamoeba histolytica</i>	cEH-P1 clone	482	GCAACTACTGTTAGTTA CCTCCAAGATATGTTTAAAC (SEQ ID NO:14)	42	25-35
	<i>Escherichia coli</i>	<i>afa</i>	750	GCTGGGCAGCAAAGTATAACTCT C CATCAAGCTGTTTGTTCGTCCGCC G (SEQ ID NO:15)	65	25

**Tabl 3: Examples of Specific Target Sequences (Cont'd)**

5	<i>Escherichia coli</i>	<i>elt</i>	275	TTACGGCGTTACTATCCTCTCTA GGTCTCGGTCAGATATGTGATTC (SEQ ID NO:16)	55	30
	<i>Escherichia coli</i>	<i>eltB</i>	195	CCTCTCTATATGCACACGGAGCTC CCAG CTATATTGTTGACTGCCCGGGACT TCGACC (SEQ ID NO:17)	54	40
	<i>Escherichia coli</i>	<i>sltIva</i>	490	TGTGTTTGCTTGTCTTCAGC ATGCGCGGGTCATGGAACGT (SEQ ID NO:18)	55	30
10	<i>Escherichia coli</i>	<i>elt</i>	750	GGCGACACATTATACCGTGC CCGAATTCTGTTATATATGTC (SEQ ID NO:19)	43	26
	<i>Escherichia coli</i>	<i>est</i>	186	TCTGTATTGTCTTTTCACC TTAATAGCACCCGGTACAAGC (SEQ ID NO:20)	43	26
	<i>Escherichia coli</i>	<i>estA</i>	175	TTTTTCTGTATTTCTTTIICII CTTIIITCAG GCAGGATTACAACAIAITTCACAG C (SEQ ID NO:21)	55	30
15	<i>Escherichia coli</i>	<i>malB</i>	595	TCGCCACACGCTGACGCTGACCA TTACATGACCTCGCTTAGTTTAC AGA (SEQ ID NO:22)	60, 65, 70	65
	<i>Escherichia coli</i>	<i>papC</i>	328	GACGGCTGTACTGCAGGGTGTGGC G ATATCCTTTCTGCAGGGATGCAAT A (SEQ ID NO:23)	65	25
	<i>Escherichia coli</i>		410	CTCCGAGAACTGGGTGCATCTTA C CGGAGGAGTAATTACAAACCTGGC A (SEQ ID NO:24)	65	25
20	<i>Escherichia coli</i>	<i>sltA</i>	140	ACCTGTAAACGAAGTTTGCG ATCTCATGCGACTACTTGAC (SEQ ID NO:25)	55	30
	<i>Escherichia coli</i>	STI (a and b)	?	TTAATAGCACCCGGTACAAGCAGG CCTGACTCTTCAAAGAGAAAATT AC (SEQ ID NO:26)	58	35
	<i>Escherichia coli</i>	<i>toxA</i>	110	CCGTATTACAGAAATCTGA GTGCATGATGAATCCAGGGT (SEQ ID NO:27)	55	35
25	<i>Escherichia coli</i>	<i>toxB (eltB)</i>	298	CAGTCTATTACAGAACTATG CCATACTGATTGCCGCAATTG (SEQ ID NO:28)	30	25
	<i>Escherichia coli</i>	<i>stx/slt-IA</i>	680	GACAGGATTTGTTAACAGG TTCCAGTTACACAATCAGGCC (SEQ ID NO:29)	55	30
	<i>Escherichia coli</i>	<i>sltI</i>	614	AACTGGATGATCTCAGTGG CTGAATCCCCCTCCATTATG (SEQ ID NO:30)	60	35

**Table 3: Examples of Specific Target Sequences (Cont'd)**

5	<i>Escherichia coli</i>	sltI	779	CCATGACAACGGACAGCAGTT CCTGTCAACTGAGCACTTTG (SEQ ID NO:31)	60	35
	<i>Escherichia coli</i>	SLT-IA	370	AAATCGCCATTCTGTTGACTACTTC T TGCCATTCTGGCAACTCGCGATGC A (SEQ ID NO:32)	60	35
	<i>Escherichia coli</i>	SLT-IIA	283	CAGTCGTCACCTACTGGTTTCATC A GGATATTCTCCCCACTCTGACACC (SEQ ID NO:33)	60	35
10	<i>Escherichia coli</i>	SLTIA, SLTIIA	224,227	TTTACGATAGACTTTTCGAC CACATATAAATTATTTGCTC (SEQ ID NO:34)	43	30
	<i>Escherichia coli</i>	slt-II (vtx2)	285	AAGAAGATGTTTATGGCGGT CACGAATCAGGTTATGCCTC (SEQ ID NO:35)	55	?
	<i>Escherichia coli</i>	vtx2	385	CATTACAGTAAAAGTGGCC GGGTGCCTCCCGGTGAGTTC (SEQ ID NO:36)	45	?
15	<i>E. coli, Salmonella, and Shigella</i>	lamB	346	CTGATCGAATGGCTGCCAGGCTCC CAACCAGACGATAGTTATCAGCA (SEQ ID NO:37)	60	30
	<i>E. coli and Shigella</i>	Putative inv	760	TAATACTCCTGAACGGCG TAGGTGTCGGCTTTTCTG (SEQ ID NO:38)	55-65	30
	<i>E. coli and Shigella</i>	stx, sltI	130	GAAGAGTCCGTGGGATTACG AGCGATGCAGCTATTAATAA (SEQ ID NO:39)	55	30
20	<i>E. coli and Shigella</i>	uidA	166	TATGGAATTCGCCGATTTT TGTTGCTCCTGCTGCGG (SEQ ID NO:40)	50	25
	<i>E. coli and Shigella</i>	uidR uidR	153 (154)	TGTTACGTCCTGTAGAAAGCCC AAAACGCTGCGCACAGCAATT (SEQ ID NO:41)	59 60	25 (30)
	<i>Shigella</i>	ail ial	320	CTGGATGGTATGGTGAGG GGAGGCCAACAATTATTTC (SEQ ID NO:42)	43 43	26 25-30
25	<i>S. dysenteriae</i>	ipaH	700	GTTCTTGACCGCCTTCCGATAC GCCGGTCAGCCACCCTC (SEQ ID NO:43)	60	35
	<i>Giardia spp.</i>	Giardin	171	AAGTGCCTCAACGAGCAGCT TTAGTGCTTTGTGACCATCGA (SEQ ID NO:44)	60	25
	<i>G. duodenalis</i>	Giardin	218	CATAACGACGCCATCGCGGCTCTC AGGAA TTTGTGAGCGCTTCTGTCTGGCA GCGCTAA (SEQ ID NO:45)	60	25

**Table 3: Examples of Specific Target Sequences (Cont'd)**

5	<i>Helicobacter pylori</i>	16S rDNA	109	CTGGAGAGACTAACGGGTGG ATTACTGACGCTGATTCTGC (SEQ ID NO:46)	60	40
	<i>Helicobacter pylori</i>	16S rRNA	~500	TGGCAATCAGCGTCAGGTAATG GCTAAGAGATCAGCCTATGTCC (SEQ ID NO:47)	55	40
	<i>Helicobacter pylori</i>	<i>ureA</i>	412	GCCAATGGTAAATTAGTT CTCCTTAATTGTTTAC (SEQ ID NO:48)	?	26
	<i>Helicobacter pylori</i>	<i>ureA</i> and <i>ureB</i>	2396	AGGAGGATGAGATGA ACTTTATTGGCTGGT (SEQ ID NO:49)	50	30
10	<i>Listeria</i> spp	16S rDNA	938	CAGCMGCCGCGGTAATWC CTCCATAAAGGTGACCCT (SEQ ID NO:50)	50	30
	<i>Listeria</i> spp	<i>iap</i>	1454	ATGAATATGAAAAAGCAAC TTATACGCGACCGAAGCCAA (SEQ ID NO:51)	50	30
15	<i>L. monocytogenes</i> and <i>L. innocua</i>	<i>iap</i>	1448	GCTACAGCTGGGATTGCGGT TTATACGCGACCGAAGCCAA (SEQ ID NO:52)	55	40
	<i>L. innocua</i>	<i>iap</i>	~870	ACTAGCACTCCAGTTGTAA TTATACGCGACCGAAGCCAA (SEQ ID NO:53)	62	30
	<i>L. invanovii</i> , <i>L. seeligeri</i> , and <i>L. welshimeri</i>	<i>iap</i>	1243	TTACTGAGGTAGCRAGC TTATACGCGACCGAAGCCAA (SEQ ID NO:54)	58	30
20	<i>L. monocytogenes</i>	16S rRNA	70	CACGTGCTACAATGGATAG AGAATAGTTTATGGGATTAG (SEQ ID NO:55)	48	40
	<i>L. monocytogenes</i>	<i>hlyA</i>	417	CATCGACGGCAACCTCGGAGA ATACAATTACCGTTCTCCACCATT C (SEQ ID NO:56)	62	30
25	<i>L. monocytogenes</i>	<i>hlyA</i>	520	AACCTATCCAGGTGCTC CGCCACACTTGAGATAT (SEQ ID NO:57)	60	35
	<i>L. monocytogenes</i>	<i>hlyA</i>	174	GCATCTGCATTCAATAAAGA TGTCACCTGCATCTCCGTGGT (SEQ ID NO:58)	56	30
	<i>L. monocytogenes</i>	<i>hlyA</i>	234	CGGAGGTTCCGCAAAGATG CCTCCAGAGTGATCGATGTT (SEQ ID NO:59)	50, 55 56	30 30
30	<i>L. monocytogenes</i>	<i>hlyA</i>	606	Sequence not reported	51	30

**Table 3: Examples of Specific Target Sequences (Cont'd)**

	<i>L. monocytogenes</i>	<i>hlyA</i>	685 <sup>c</sup>	CCTAAGACGCCAATCGAA AAGCGCTTGCAACTGCTC (SEQ ID NO: 60)	50	30
5	<i>L. monocytogenes</i>	<i>hlyA</i>	702	CCTAAGACGCCAATCGAA AAGCGCTTGCAACTGCTC (SEQ ID NO: 61)	50	30
	<i>L. monocytogenes</i>	<i>hlyA</i>	234	ATTGCGAAATTTGGTACAGC ACTTGAGATATATGCAGGAG (SEQ ID NO: 62)	55	30
	<i>L. monocytogenes</i>	downstream from <i>hlyA</i>	?	Sequences not reported	55	30
10	<i>L. monocytogenes</i>	<i>iap (msp)</i>	131	ACAAGCTGCACCTGTTGCAG TGACAGCGTGTGTAGTAGCA (SEQ ID NO: 63)	55	30
	<i>L. monocytogenes</i>	<i>iap</i>	544	CAAGCAACTACACCTGCGCC GAACCTTGTTAGCATTCTGT (SEQ ID NO: 64)	50	30
15	<i>L. monocytogenes</i>	<i>msp (iap)</i>	593	ACAAGCTGCACCTGTTGCAG GAACCTTGTTAGCATTCTGT (SEQ ID NO: 65)	50	30
	<i>L. monocytogenes</i>	<i>iap</i>	287	CGAATCTAACGGCTGGCACA GCCCAAATAGTGTACCGCT (SEQ ID NO: 66)	50	30
	<i>L. monocytogenes</i>	<i>iap</i>	380+/- 40 <sup>d</sup>	CAAAGTCTAACACAGCTACT GCACTTGAATTGCTGTTATTG (SEQ ID NO: 67)	65	35
20	<i>L. monocytogenes</i>	Dth-18	326	CCGGGAGCTGCTAAAGCGGT GCCAAACACCGAAATACC (SEQ ID NO: 68)	54	30
	<i>L. monocytogenes</i>	Dth-18	122	GAAGCACCTTTTGACGAAGC GCTGGTGCTACAGGTGTTTC (SEQ ID NO: 69)	54	30
25	<i>L. monocytogenes</i>	<i>lmaA</i>	257	AACAAGGTCTAACTGTAAAC ACTATAGTCAGCTACAATTG (SEQ ID NO: 70)	55	30
	<i>Salmonella</i>	DNA rep. ori.	163	TTATTAGGATCGCGCCAGGC AAAGAATAACCGTTGTTTAC (SEQ ID NO: 71)	50	35
	<i>Staphylococcus aureus</i>	<i>entA</i>	120	TTGGAACCGGTTAAACGAA GAACCTTCCCATCAAAAACA (SEQ ID NO: 72)	55	?
30	<i>Staphylococcus aureus</i>	<i>entB</i>	478	TCGCATCAAACGTACAAACG GCAGGTACTCTATAAGTGCC (SEQ ID NO: 73)	55	?

**Table 3: Examples of Specific Target Sequences (Cont'd)**

5	<i>Staphylococcus aureus</i>	<i>entB</i>	593	GAGAGTCAACCAGATCCTAAACCA G ATACCAAAGCTATTCTCATTTTC T (SEQ ID NO:74)	50	50
	<i>Staphylococcus aureus</i>	<i>entC1</i>	801	ATGAATAAGAGTCGATTATTTCAT T TTATCCATTCTTTGTTGTAAGGTG G (SEQ ID NO:75)	50	50
	<i>Staphylococcus aureus</i>	<i>entC1</i>	631	ACACCCAAAGTATTAGCAGAGAG CC CCTGGTGCAGGCATCATATCATA CC (SEQ ID NO:76)	50	50
10	<i>Staphylococcus aureus</i>	<i>entC1</i>	257	GACATAAAAGCTAGGAATTT AAATCGGATTAACATTATCC (SEQ ID NO:77)	55	?
	<i>Staphylococcus aureus</i>	<i>entD</i>	317	CTAGTTTGGTAATATCTCCT TAATGCTATATCTTATAGGG (SEQ ID NO:78)	55	?
	<i>Staphylococcus aureus</i>	<i>entE</i>	170	TAGATAAAGTTAAAACAAGC TAACCTACCGTGGACCCTTC (SEQ ID NO:79)	55	?
15	<i>Staphylococcus aureus</i>	<i>ist</i>	186	TTCATATTGTAAAAGTGTCAG ACCCCACT TACTAATGAATTTTTTTATCGTA AGCCCTT (SEQ ID NO:80)	55	40
	<i>Staphylococcus aureus</i>	<i>ist</i>	350	ATGGCAGCATCAGCTTGATA TTCCAATAACCACCGTTT (SEQ ID NO:81)	55	?
	<i>Staphylococcus aureus</i>	<i>eat</i>	119	CTAGTGCATTTGTTATTCAA TGCATTGACACCATAGTACT (SEQ ID NO:82)	55	?
20	<i>Staphylococcus aureus</i>	<i>eth</i>	200	ACGGCTATATACATTCAATT TCCATCGATAATATACCTAA (SEQ ID NO:83)	55	?
	<i>Staphylococcus aureus</i>	<i>nuc</i>	450	AGTATATAGTGCAACTTCAACTA AA ATCAGCGTTGTCTTCGCTCCAAA TA (SEQ ID NO:84)	50	50
	<i>Staphylococcus aureus</i>	<i>nuc</i>	-270	GCGATTGATGGTGATACGGTT AGCCAAGCCTTGACGAACCTAAAG C (SEQ ID NO:85)	55	37
25	<i>Vibrio cholerae</i>	<i>ctx</i>	302	CTCAGACGGGATTTGTTAGGCAC G TCTATCTCTGTAGCCCCCTATTAC G (SEQ ID NO:86)	60	30, 40
	<i>Vibrio cholerae</i>	<i>ctx</i>	384	CGGGCAGATTCTAGACCTTC GCACCCCAAATAGAACTCGA (SEQ ID NO:87)	62	25



**Table 3: Examples of Specific Target Sequences (Cont'd)**

	<i>Vibrio cholerae</i>	<i>ctxA</i>	564	CGGGCAGATTCTAGACCTCCTG CGATGATCTTGGAGCATTCCCAC (SEQ ID NO:88)	60	55
	<i>Vibrio cholerae</i>	<i>ctxAB</i>	777	TGAAATAAAGCAGTCAGGTGGTG ATTCTGCACACAAATCAG GGTATTCTGCACACAAATCAG (SEQ ID NO:89)	55	25-35
	<i>Vibrio cholerae</i>	<i>luxA</i>	350	GGAAGCTTCCAATGATTCTAAGC TGGAT GGGAATTCTCAGGCGTCCCTACT GGGTT (SEQ ID NO:90)	55	25
5	<i>V. parahaemolyticus</i>	<i>idh</i>	648	CTGTCCCTTTTCTGCCCCCG GCTCTTAGCTGCGGCGGTGGT (SEQ ID NO:91)	62	25
	<i>V. vulnificus</i>	<i>Cytolysin</i>	388	CGCCGCTCACTGGGGCACTGGCT G CCAGCCGTTAAGCGAACCACCCG C (SEQ ID NO:92)	65	40, 50
	<i>V. vulnificus</i>	<i>Cytolysin</i>	519	CCGCGGTACAGGTTGGCGCA CGCCACCCACTTTCGGGCC (SEQ ID NO:93)	67-69	30
10	<i>Yersinia enterocolitica</i>	<i>ail</i>	273	GAACTCGATGATAACTGGG GCAATTCAACCCACTTCAA (SEQ ID NO:94)	65	35
	<i>Yersinia enterocolitica</i>	<i>ail</i>	170	ACTCGATGATAACTGGGGAG CCCCAGTAATCCATAAAGG (SEQ ID NO:95)	55	25, 30
15	<i>Yersinia enterocolitica</i>	<i>ail</i>	425	TTAATGTGTACGCTGCGAGTG GGAGTATTCATATGAAGCGTC (SEQ ID NO:96)	57	35
	<i>Yersinia enterocolitica</i>	<i>inv</i>	359	CTATTGGTTATGCGCAAAGC TGGAAGTGGGTTGAATTGCA (SEQ ID NO:97)	?	?
	<i>Yersinia enterocolitica</i>	<i>yst</i>	163	AATGCTGTCTTCATTGGAGC GCAACATACATCACAGCAATC (SEQ ID NO:98)	60	35
20	<i>Yersinia enterocolitica</i>	<i>yst</i>		AAAGATATTTTGTCTTGT GCAGCCAGCACACGCGGG (SEQ ID NO:99)	43	30
	<i>Y. pseudotuberculosis</i>	<i>inv</i>	295	TAAGGGTACTATCGAGGCGGA CGTGAAATTAACCGTCACACT (SEQ ID NO:100)	55	25, 30
25	<i>Yersinia virulence plasmid</i>	<i>virF</i>	590	TCATGGCAGAACAGCAGTCAG ACTCATCTTACCATTAAGAAG (SEQ ID NO:101)	?	?

**Table 3: Examples of Specific Target Sequences (Cont'd)**

<i>Yersinia virulence plasmid</i>	<i>virF</i>	590	CATGGCAGAACAGCAGTCAG ACTCATCTTACCATTAAGAAG (SEQ ID NO:102)	57	35
<i>Norwalk virus</i>	<i>Near 3' end</i>	456	ATTGAGAGCCTCCGCGTG GGTGGCGAAGCGGCCCTC (SEQ ID NO:103)	49	30-40
<i>Norwalk virus</i>	<i>pol</i>	470	CTTGTTGGTTTGAGGCCATAT ATAAAAGTTGGCATGAACA (SEQ ID NO:104)	55	40
<i>Norwalk virus</i>	<i>pol</i>	260	CAAATTATGACAGAATCCTTC GAGAAATATGACATGGATTGC (SEQ ID NO:105)	55	40
<i>Norwalk virus</i>	<i>IP</i>	224	CACCACCATAAACAGGCTG AGCCTGATAGAGCATTCTTT (SEQ ID NO:106)	50	40
<i>Rotavirus</i>	<i>vp7</i>	1036	GGCTTTAAAAGAGAGAATTTCCGT CTGG GGTCACATCATACAATTCCTAATC TAAG (SEQ ID NO:107)	42	25
<i>Rotavirus</i>	<i>Gottfried gene 4 seg.</i>	?	CCATATCAGCCAACGAGT TTACTACTTCTACATCAGGT (SEQ ID NO:108)	42	30
<i>Rotavirus</i>	<i>OSU gene 4 seg.</i>	?	CATACCAACCAACCACTTTC TGATGTCATATTTACTGTGT (SEQ ID NO:109)	42	30
<i>Rotavirus (group A)</i>	<i>near 5' end</i>	259	GGCTTTTAAACGAAGTCTTC TCAACAATGCGTCTAAGTTCACAG (SEQ ID NO:110)	55	30
<i>Hepatitis A virus</i>	<i>VP1</i>	302	TCCCAGAGCTCCATTGAA CATTATTTTCATGCTCCTCAG (SEQ ID NO:111)	37	30
<i>He</i>	<i>VP3</i>	207	ACAGGTATACAAAGTCAG CTCCAGAATGATCTCC (SEQ ID NO:112)	37	30
<i>Hepatitis A virus</i>	<i>VP3</i>	207	CTCCAGAATCATCTCC ACAGGTATACAAAGTCAG (SEQ ID NO:113)	49	40
	?	?	Sequences not reported	?	30
<i>Enteroviruses</i>	<i>Near 5' end</i>	154	CCTCCGGCCCTGAATGCGGCTAA T TAACAGTGGATTCGTCGGT (SEQ ID NO:114)	50	25, 50

Species specific primers and nucleic acid target sequences for *C. jejuni* are provided in

Table 4 below. See, Day et al., Appl. Environ. Microbiol.: 1019 (March 1997).

**Table 4:** Further Species Specific Sequences

5	Organism	Primers (5' to 3')	Target Sequence (5' TO 3')
10	<i>C. Jejuni</i>	AGTCAGCCAC (SEQ. ID NO: 126) AATCGGGATG (SEQ. ID NO: 128) AGGGGTCTTG (SEQ. ID NO: 130) AGCCAGCGAA (SEQ. ID NO: 132) GAAACGGGTG (SEQ. ID NO: 134) GTGACGTAGG (SEQ. ID NO: 136)	ATCGGGCTGTTATGATGATA (SEQ. ID NO: 127) ATCACTGGGGGAGCTAATAT (SEQ. ID NO: 129) TAAGGTAAAGTTGTTGTGAATC (SEQ. ID NO: 131) CATATCCAGAGCCTCTGGAT (SEQ. ID NO: 133) GTAGCCTCTTCATCGTCGTCTAA (SEQ. ID NO: 135) CACCCGCTTAACGCCAAGA (SEQ. ID NO: 137)

The construction of the DNA beacon molecule, PCR reaction conditions, and sample preparation are detailed below. Briefly, generally known PCR reaction conditions are suitable. For example, a master pre-mix for a standard 50 µl reaction, although the volume may be reduced to as small as a 10 µl per reaction, may be as follows. 24 µl water free of nucleases, proteases, heavy metals and DNA/RNA may be mixed with 5 µl of a 2% PVPP aqueous solution (polyvinylpolypyrrolidone), 5 µl of 10 x PCR buffer, e.g. 100 mM tris-HCl, pH 8.3, 500 mM KCl, 5 µl of 25 mM MgCl<sub>2</sub>, 1.25 µl 50 ng/µl of each primer, 1 µl 10 mM of each dNTP, 2.5 µl of enzyme, e.g. AmpliTaq LD at a 1:5 dilution and 2 µl of DNA template preparation, as described below. The cycling parameters may be 30 sec. at 95°C, 30 sec. at 58°C and 60 sec. at 74°C. This routine may be repeated 30-50 times depending on the sensitivity needed, which is generally empirically determined by the type of sample and sample preparation without undue experimentation.

The preparation of the sample is adapted empirically based on whether or not there are interfering substances in the sample, and what type of substances they are, as is known in the art. Once the proper sample preparation is attained, DNA extraction may be attained, for example, by simple boiling for 10 min prior to utilization of the sample as DNA template or target in the reaction. For example, to test for sterility of shelf stable commercially sterile pudding, the pudding may be diluted in ultrapure water 1:30 and boiled for 10 min before adding to the PCR pre-mix.

The fluorescent beacon is constructed by assembly of the separate parts, that is the nucleic acid and the fluorophore and quencher are synthesized separately and then bound by means of linkers, e.g. amino linkers, as is known in the art.

A large variety of microorganisms are involved in the contamination of, for example,

food, cosmetics, medical supplies and fluids, blood products, intravenous solutions, and the like, among other commercial products. Examples of these are *E. coli*, *Salmonella*, *Bacillus*, *Clostridium*, *Listeria*, *Pseudomonas*, and many others including those listed above, all of which are promptly detected by the present technology, which is suitable for the standardized testing of products samples to determine their suitability for administration to or use in humans or animals. A necessarily incomplete list of microorganisms commonly present in foodstuffs is listed by Hill W. E., in Critical Rev. Food Sci. and Nutrition 36 (1 & 2): 123 (1996), along with other universal and specific bacterial primers and PCR cycling temperature pairs, the relevant pages thereof being incorporated herein by reference, e.g. pp. 140-146, among others. Other universal and specific sequences of internal DNA segments plus DNA arms including the 5' and 3' termini of the beacon are also provided by Schofield et al., Appl. and Environ. Microbiol. 63(3): 1143 (1997), the relevant sections of which are incorporated herein by reference, e.g., Tables 1 and 2, and the section on Cultivation of Bacteria, among others.

The reaction conditions for nucleic acid amplification may be custom tailored for the specific primers employed, and an artisan would know how to easily and quickly determine suitable conditions for specific applications. When the sample to be analyzed is suspected of containing a sufficient quantity of bacteria, generally a nucleic acid sequence about 100 and up to about 500 nucleotide long, and preferably up to about 200 base pairs long or even longer, may be amplified, and a polynucleotide DNA (or fluorescent beacon probe) designed so that it hybridizes to a portion of the DNA fragment. However, longer and smaller segments of nucleotide may also be utilized. When hybridized to a target, the amount of fluorescence emitted by polynucleotide plus fluorogenic reagents (or by a fluorescent beacon) is proportional to the amount of amplified nucleic acid present in the sample. The PCR conditions are generally set up so that the expected nucleic acid product comprises about 100 about 400 base pairs and contains, among other sequences, the target sequence for the complementary polynucleotide DNA internal segment (probe). However, other lengths are also suitable. The PCR may be conducted for as many cycles as necessary to obtain enough nucleic acid to bind the DNA beacon and obtain a linear fluorescent signal. In most cases up to about 30 PCR cycles are sufficient to produce enough nucleic acid target for the the DNA polynucleotide (or beacon) to bind to, and to produce sufficient fluorescence for observation with the naked eye after gel electrophoresis. A specialized instrument or fluorometer may be utilized for the measurement of fluorescence during the course of the reaction, and the display of typical response curves.

A sample may be assayed side-by-side with proper positive and negative controls, as is known in the art, and whether or not the sample has bacterial contamination may be conclusively determined in as short a time as 10 minutes. Positive controls may be conducted by following the same steps performed on the test sample but by adding to the medium a target  
5 nucleic acid sequence, e.g. naked DNA or bacteria, viruses, and the like. Negative controls are conducted in a manner similar to that of the test sample but contain no target nucleic acid.

This invention was described above in relation to the universal detection of microorganisms and/or viruses, and for the specific detection of a class of microorganism(s), e.g. bacteria, as well as for implementation with one beacon and two DNA primers. The  
10 invention is, however, not limited to either the described sequences, nor the number of polynucleotides (or fluorescent beacons) and primers. In fact, any combination of primer and polynucleotide (or beacon) sequences that are specific or universal for microorganisms and viruses, including yeast, molds, and the like, may be substituted and/or added to the nucleic acid target subjected to PCR. Examples of nucleic acid targets which may be utilized are those  
15 specific for a microorganism, or genus or class of microorganisms, such as a nucleic acid segment imparting a virulence trait, DNA sequences that are order-, class-, strain- and/or species-specific, such as those based on immunological identity, e.g. antigenic proteins or flagella. A partial list of DNA target probe sequences and their corresponding primers is provided by Hill, E. W. et al., Critical Rev. Food Sci. & Nutrition 36(182):123 (1996), the  
20 relevant text of which is incorporated herein by reference.

Similarly, nucleic acid targets and primers may also be utilized to design polynucleotides (or fluorescent beacons) which specifically hybridize to a single species of virus or microorganism, bacteria such as *E. coli*, *Salmonella*, *Listeria*, etc., viruses such as Rotavirus, Influenza, etc., among others. Thus, the assay and the kit of the invention may also  
25 be targeted to the determination of specific microorganisms and/or viruses. The kit of this invention along with a instructions to custom tailor and perform PCR may be applied for the intended use in conjunction with any commercially available thermocycler. One embodiment of this invention, therefore, provides a kit which contains a pre-mixed composition comprising specific nucleic acid primers and beacon probes, which may carry a fluorescent molecule.  
30 Alternatively, the fluorogenic tag or beacon may be provided separately for attachment prior to use. The reaction mixture (pre-mix) is intended for direct mixing with a sample, and for practicing the present assay detecting the presence of microorganisms and viruses in as short a time as about 10 minutes after sample preparation. The kit of the invention may be pre-

packaged in bulk for aliquoting by the user, or in unit form, as lyophilized or vitriculated form, in reaction vessels to ensure the stability of the components during shipping and storage. When in lyophilized or vitriculated form, the user may simply rehydrate the reaction ingredients in the vessel containing them or upon aliquoting, with a predetermined amount of sample preparation and conduct the detection step by placing the vessel in the fluorimeter.

The user of the kit of this invention need not be technically skilled to perform the assay provided here, to obtain accurate results. The present inventor has discovered and designed DNA segments for the universal detection of microorganisms, e.g. bacteria. Other DNA segments complementary to known nucleic acid sequences which have specificity for one or the other microorganism or virus may be designed and amplified by known PCR reactions for use with the present kit and assay. PCR reaction conditions are widely known in the art, and known and to-be-discovered universal and specific nucleic acid sequences, and primers, may be utilized with this technology. See, Hill (1996), supra.

The method and kit of the invention find numerous suitable applications, for example, in the food and cosmetic industry, as well as for the clinical diagnostic industry, e.g. for the detection of general or specific microorganisms and viruses, with simple instrumentation, and unskilled technicians.

### EXAMPLES

#### Example 1: Typical PCR Reaction Components & Conditions

A master pre-mix for a standard 50  $\mu$ l reaction (may be reduced to as small as a 10  $\mu$ l reaction) is prepared as follows.

24  $\mu$ l ultra purified water free of nucleases, proteases, heavy metals and DNA/RNA are mixed with 5  $\mu$ l of a 2% solution of PVPP (polyvinylpolypyrrolidone) in ultra purified water, 5  $\mu$ l of 10 x PCR buffer (100 mM tris-HCl, pH 8.3, 500 mM KCl in ultrapure water), 5  $\mu$ l of  $MgCl_2$  (25 mM solution in ultrapure water), 1.25  $\mu$ l of each primer (2 solutions at 50 ng/ $\mu$ l in ultrapure water), 1  $\mu$ l of each dNTP (4 solutions at 10 mM in ultrapure water), 2.5  $\mu$ l of enzyme (AmpliTaq LD at a 1:5 dilution in ultrapure water, Perkin-Elmer) and 2  $\mu$ l of DNA template preparation as described below.

The mixture is then cycled under the following conditions: 30 sec at 95°C, 30 sec at 58°C and 60 sec at 74°C. Each cycle is repeated 30-50 times depending on the sensitivity needed (determined by sample type and sample preparation, empirically).

**Example 2: Sample Preparation**

The conditions for the preparation of the sample are determined empirically based on the type and amount of the expected interfering substances to be present in the sample. However, once the proper blend of sample is finalized, a simple boil DNA extraction for 10 min is conducted prior to utilizing the sample as DNA template in the reaction.

To test for sterility of shelf stable commercially sterile pudding, for example, the pudding is diluted in ultrapure water 1:30 times, and boiled for 10 min before adding the pudding sample to a PCR pre-mix.

**Example 3: Beacon Synthesis**

Custom oligonucleotides having specified nucleotide sequences, a 5'-methoxytrityl protected C<sub>6</sub> thio group, and a 3'-C<sub>7</sub> amino linker group were utilized as starting materials for the synthesis. The synthetic sequence first converts the amine group of the oligomer to its dabcyll amide derivative, removes the methoxytrityl protective group, and converts the resulting thiol to edans or fluorescein thioether derivatives. Tyagi and Kramer's general synthetic strategy is employed by following Tyagi's supplemental procedures. See, Tyagi, S. And Kramer, R.R., Molecular Beacons: Probes that Fluoresce upon Hybridization. Nature Biotechnology 14:303-308 (1996); modified as described here under the advise of Tyagi, S., Preparation of Molecular Beacons, Private Communication (1996).

Tyagi's procedure was modified for the present work in several ways, most significantly by changing reaction conditions during attachment of the edans or fluorescein groups, deletion of the precipitation step during the purification of the dabcyllated intermediate and deletion of both HPLC and precipitation steps during purification of the final beacon product. In our hands, the precipitation of the dabcyll intermediate and the beacon product from ethanolic solution either did not occur or resulted in substantial losses of material. Under the conditions described here, conversion of the dabcyll intermediate to the beacon proceeded cleanly and, in many cases, HPLC purification of the product afforded little benefit.

**DETAILED PREPARATION OF CODE 23 S-B'-FLUORESCIEIN BEACON****Example 4: Reconstitution of DNA Oligomer**

The DNA oligomer is a lyophilized solid obtained from Midland Certified Reagent Co. (Lot 021098-175, 2008 nmoles). It was dissolved in 3.0 mL 0.1 M carbonate buffer (pH = 8.5)

and 500  $\mu$ L used for the following dabcylation reaction.

**Example 5: Conversion of Oligomer to DabcyL Derivative**

Aliquots (10  $\mu$ L) of a saturated solution of se-dabcyL (Molecular Probes, 4-((4-dimethylamino)phenyl)azo)benzoic acid, succinimidyl ester) in dimethylformamide (DMF) were added at approximately 30 min intervals to the stirred oligomer. The se-dabcyL reagent was made by mixing 12.2 mg se-dabcyL with 110  $\mu$ L DMF. A total of 10 aliquots are added. The reaction was allowed to proceed at ambient temperature for about 29 hours.

**Example 6: Purification of the DabcyL Intermediate**

The reaction mixture was transferred to a small centrifuge tube, centrifuged for 4 min. at 14000 rpm, and the supernatant transferred to a second centrifuge vial. Residual material in the reaction flask was washed with a total of 400  $\mu$ L of 0.01M triethylammonium acetate (TEAA, pH = 6.5), and the washings used to resuspend the solids from the first centrifugation. After centrifugation, the supernatant was added to that from the first centrifugation. The total volume of the combined supernatants was reduced to 500  $\mu$ L by vacuum evaporation (Speed Vac). Unreacted se-dabcyL and other relatively small compounds were separated from the dabcyL derivative and other oligomeric species by gel permeation chromatography using an NAP-5 (Pharmacia) column and 0.01M TEAA mobile phase. See below, for specific procedure used for a NAP-5 column). The fraction containing the higher molecular weight materials was concentrated to slightly less than 500  $\mu$ L. DabcyL intermediate was then separated from other oligomers via reversed phase HPLC (Vydac C18 column, see note 2 for procedure) of two approximately 250  $\mu$ L injections.

Under the HPLC conditions employed, the present dabcyL intermediate had a retention time of about 28-30 min and yields yellowish fractions. Several 1 mL fractions in this region were pooled and concentrated to 500  $\mu$ L. The dabcyL intermediate was desalted by NAP-5 column chromatography using 0.01M TEAA mobile phase and the resulting 1 mL eluate concentrated to about 250  $\mu$ L.

**Example 7: Conversion of DabcyL Intermediate to Fluorescein Beacon**

After removing a 5  $\mu$ L sample (for analysis by HPLC, note 3), 10  $\mu$ L of a freshly prepared aqueous silver nitrate (0.27 g  $\text{AgNO}_3$ /10 mL, about a 5-fold molar excess of  $\text{AgNO}_3$  over DNA oligomer, solution was added and the mixture allowed to react for 30 min at room



temperature with intermittent mixing. See, Connolly, B.A., Chemical Synthesis of Oligonucleotides Containing a Free Sulfhydryl Group and Subsequent Attachment of Thiol Specific Probes, Nucleic Acids Research 13 (12): 4485-4503 (1985).

15  $\mu\text{L}$  of freshly prepared aqueous dithiothreitol solution (0.23 dithiothreitol/10 mL, about a 7-fold molar excess over DNA oligomer) were added and reaction proceeded for 5 min at ambient temperatures with occasional stirring. The reaction mixture was centrifuged, and the supernatant transferred to a fresh reaction vial. Another 5  $\mu\text{L}$  analytical sample was removed. The pH of the remaining supernatant was then adjusted to about 9.0 by treating with an equal volume (about 265  $\mu\text{L}$ ) of 0.2M carbonate (pH = 9.0). To this, 25  $\mu\text{L}$  of a freshly prepared  
10 solution of 5-IAF in dimethylsulfoxide (DMSO) was added. This solution contained about 10.9 mg 5-IAF (Molecular Probes, 5-IAF iodoacetamidofluorescein) in 125  $\mu\text{L}$  DMSO. This is about a 13.5-fold excess of 5-IAF over oligomer and about a 2.5-fold molar excess of 5-IAF over total sulfhydryl groups (deprotected DNA oligomer and dithiothreitol). After brief  
15 mixing, reaction was allowed to proceed for about 3.5 days at refrigerator temperatures (about 5°C).

**Example 8: Purification of Fluorescein Beacon**

The volume of the reaction mixture was reduced to 500  $\mu\text{L}$  and the crude reaction mixture chromatographed on an NAP-5 column using 2 mM Tris buffer (pH = 8.3, Trizma, Sigma). The higher molecular weight fraction was concentrated to 500  $\mu\text{L}$  and again purified  
20 on an NAP-5 column with Tris mobile phase. The volume of the final preparation was adjusted to 500  $\mu\text{L}$ .

**Example 9: Fluorescein Beacon Yield**

The concentration on the final beacon product was determined spectrophotometrically (absorbance at 260 nm) after correcting the observed reading for the spectral contribution of  
25 the dabcyI and fluorescein groups.

$$\text{Abs 260 (corr)} = \text{Abs 260 (obs)}/1.14$$

The yield from this preparation was about 31% based on amount of starting DNA oligomer.

**Exempl 10: NAP-5 Gel Permeation Chromatography**

5 A NAP-5 column was equilibrated to the particular mobile phase and used according to the manufacturer's directions. This entails allowing the excess liquid present with the commercial column to elute, equilibration with about 10 mL of mobile phase, and loading the sample (volume 500  $\mu$ l max). During loading, all of the liquid phase was allowed to enter the column bed and eluting material is discarded. Subsequently, a small centrifuge vial was positioned to collect eluant and 1.0 mL of mobile phase was added. The approximately 1.0 mL eluting contains most of the desired DNA oligomeric materials.

**Example 11: Reversed Phase HPLC**

10 A Waters instrument equipped with dual pumps, a solvent programmer (Waters 600E System Controller), Vydac C18 RP column, UV-Vis detector (Waters 486 Tunable Absorbance Detector), A/D converter (PE Nelson 900 Series Interface), computer with associated chromatographic software (PE Nelson Turbochrom), and manual injector was used. The column flow rate used was 1.0 mL/min, and the detection wavelength was usually 260  
15 (sometimes 490) nm. A linear solvent gradient changed solvent composition from 100% A to 100% B over a 40 min period. Solvent A was an 80:20 (v/v) mixture of 0.1M TEAA (pH = 6.5) and acetonitrile; solvent B a 30:70 mixture of 0.1M TEAA and acetonitrile.

**Example 12: Analytical HPLC**

20 The composition of the starting material, the final product and intermediate preparations (both the protected and deprotected dabcyI species) were monitored by HPLC under conditions identical to those described below but using 20  $\mu$ L injections. The aliquots noted above were concentrated enough that, after dilution about 10-fold (usually in 0.1M TEAA or 2mM Tris buffers), adequate responses were detected.

**Edans**

25 The edans beacon referred to above results from reacting the deprotected sulfhydryl group with 1,5-IAEDANS (Molecular Probes, 1,5-IAEDANS stands for 5-(((2-iodoacetyl)amino)ethyl)amino)naphthalene-1 sulfonic acid).

**References**

1. Tyagi, S. And Kramer, R.R., 1996. Molecular Beacons: Probes that Fluoresce upon

Hybridization. Nature Biotechnology 14:303-308.

2. Tyagi, S., 1996. Preparation of Molecular Beacons. Private communication.

3. Connolly, B.A. 1985. Chemical Synthesis of Oligonucleotides Containing a Free Sulfhydryl Group and Subsequent Attachment of Thiol Specific Probes. Nucleic Acids

5 Research 13(12): 4485-4503.

### CLAIMS

1. An in vitro method of detecting the presence of bacteria in a sample, comprising the steps of:

(a) forming a polymerase chain reaction mixture by combining (1) a pre-determined volume of a sample to be tested for the presence of a nucleic acid sequence comprising 5'-TAGAAGC-3', (2) known amounts of a first primer comprising 5'-GCTAAGGTCCCAAAGT-3' and a second primer comprising 5'-AGAACGCTCTCCTACC-3', and (3) polymerase chain reaction reagents;

(b) forming a polymerase chain reaction product by cycling the polymerase chain reaction mixture under conditions effective to amplify the nucleic acid sequence, if present, to replicate and attain about 0.25 to about 10,000  $\mu\text{g}$  nucleotide product/ $\mu\text{l}$  mixture;

(c) adding a probe containing DNA comprising 5'-GGTGGCTGCTTCTAAGCCACC-3' to the polymerase chain reaction mixture or to the polymerase chain reaction product to cause the DNA to hybridize with the nucleic acid sequence, if present, and change the conformation of the probe; and

(d) determining whether or not bacteria is present in the sample by detecting the conformational change of the probe, the conformational change indicating the presence of bacteria in the sample.

2. The method in accordance with claim 1, wherein a change in conformation of the probe is detected by a fluorescent detection system.

3. The method in accordance with claim 2, wherein a change in conformation results in fluorescence of the probe.

4. The method in accordance with claim 1, further comprising one end of the probe being operatively connected to a fluorogen and the other end of the probe being operatively connected to a quencher and wherein a change in conformation of the probe is detected by a fluorescent detection system.

5. The method in accordance with claim 4, further comprising a first linker and a second linker operatively connecting the ends of the probe to the fluorogen and the

quencher, respectively.

6. The method in accordance with claim 5, wherein the linkers are comprised of C<sub>6</sub>-thio or C<sub>7</sub>-amino linkers.

7. The method of claim 6, wherein each linker comprises about 1 to about 10 carbon atoms.

8. The method in accordance with claim 4, wherein the probe is added to the polymerization reaction mixture.

9. The method in accordance with claim 4, wherein the probe is added to the polymerization reaction product.

10. The method in accordance with claim 4, wherein the fluorogen is fluorescein the quencher is dabcyI.

11. The method in accordance with claim 1, wherein a fluorogen is operatively connected to one end of the first or second primer; and further comprising quenching any primer not bound to the polymerase chain reaction product.

12. The method in accordance with claim 11, wherein any primer not bound to the upstream or the downstream sequences in the polymerase chain reaction product is quenched by adding to the polymerase chain reaction product an oligonucleotide that is complimentary to the primer and has a quencher operatively attached to said oligonucleotide.

13. The method in accordance with claim 11, wherein the fluorogen is operatively connected to the 5' end of the first primer or the second primer and any primer that is not bound to the upstream or the downstream sequences in the polymerase chain reaction product is quenched by adding to the polymerase chain reaction product an oligonucleotide that is complimentary to the primer and has a quencher operatively attached to the 3' end of said oligonucleotide.

14. The method in accordance with claim 11, wherein the fluorogen is fluorescein and the quencher is dabcyl.

15. The method in accordance with claim 1, wherein the cycling is conducted about 1 to 30 times at alternating temperatures of about 95 to about 58° C: about 58 to about 74° C or 74:95° C.

16. The method in accordance with claim 1, wherein the probe is added at a concentration about 0.6 to about 1.6 with respect to the concentrations of the primers.

17. The method in accordance with claim 1, wherein the polymerase chain reaction reagents include an enzyme for degrading double stranded nucleotide products.

18. The method in accordance with claim 1 further comprising adding to the polymerase chain reaction product single-stranded DNA that is complementary to at least a portion of the nucleic acid sequence and causing the DNA to hybridize with at least a portion of the nucleic acid sequence, if present, to form double stranded nucleic acid; and determining whether or not the nucleic acid sequence is present in the polymerase chain reaction product by detecting any double stranded DNA.

19. The method in accordance with claim 18 wherein the formation of double stranded nucleic acid is detected by a fluorescent detection system, and wherein when no universal or specific nucleic acid is present and replicated by the polymerase chain reaction in the sample the internal DNA segment remains single stranded and no fluorescence is detected, and when there is microbial or viral nucleic acid present and replicated by the polymerase chain reaction, its microbial and/or nucleic acid sequence is double stranded and operatively bound to the fluorogenic agent fluoresces.

20. The method of claim 19, wherein the nucleotide products in the polymerase reaction product are separated prior to the fluorescence detection.

21. The method of claim 20, wherein the nucleotide products are separated by

electrophoresis.

22. The method of claim 21, further comprising staining the nucleotide products with a DNA stain that operatively binds to double stranded nucleic acid, but not to single stranded nucleic acid.

23. The method in accordance with claim 22 wherein the DNA stain is ethidium bromide and fluorescence is detected under ultraviolet light.

24. A polynucleotide consisting essentially of 5'-GCTAAGGTCCCAAAGT-3'.

25. The polynucleotide in accordance with claim 24, further comprising a fluorogen operatively connected at the 5'-end.

26. The polynucleotide in accordance with claim 25, wherein the fluorogen is fluorescein, pico green, or edans.

27. A polynucleotide consisting essentially of 5'-AGAACGCTCTCCTACC-3'.

28. The polynucleotide in accordance with claim 27, further comprising a fluorogen operatively connected at the 5'-end.

29. The polynucleotide in accordance with claim 28, wherein the fluorogen is fluorescein, pico green, or edans.

30. A polynucleotide consisting essentially of 3'-CGATTCCAGGGTTTCA-5'.

31. The polynucleotide in accordance with claim 30, further comprising a quencher operatively connected at the 3'-end.

32. The polynucleotide in accordance with claim 31, wherein the quencher is dabcyI.

33. A polynucleotide consisting essentially of 3'-TCTTGCGAGAGGATGG-5'.
34. The polynucleotide in accordance with claim 33, further comprising a quencher operatively connected at the 3'-end.
35. The polynucleotide in accordance with claim 34, wherein the quencher is dabcyl.
36. A polynucleotide consisting essentially of 3'-TAGAAGC-5' or 3'-ATCTTCG-5'.
37. A polynucleotide consisting essentially of 3'-GGTGGCT TAGAAGC AGCCACC-5', 3'-GCTGGCTATCTTCGAGCCACC-5', 3'-CCACCGAATCTTCG TCGGTGG-5', or 3'-CCACCGATAGAAGCTCGGTGG-5'.
38. The polynucleotide in accordance with claim 37, further comprising a fluorophore operatively attached at the 5'-end and a quencher operatively attached at the 3'-end.
39. The polynucleotide in accordance with claim 38, wherein the fluorophore is fluoresceine and the quencher is dabcyl.
40. The polynucleotide in accordance with claim 38, further comprising a first linker and a second linker operatively connecting the termini of the arm segments to the fluorogen and the quencher, respectively.
41. The polynucleotide in accordance with claim 40, wherein the linkers are comprised of C<sub>6</sub>-thio or C<sub>7</sub>-amino linkers.
42. The polynucleotide of claim 41, wherein each linker comprises about 1 to about 10 carbon atoms.



43. A kit for detecting a microbe or a virus, comprising in separate containers, primers selected from the group consisting of those binding to sequences up-stream and down-stream to a universal or specific nucleic acid sequence indicative of a microbe or a virus;

a polynucleotide comprising a DNA internal segment being hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence, a first and a second DNA arm segment adjoining the internal segment, each of the DNA arm segments comprising nucleotide sequences hybridizably complementary to one another; and instructions for use of the kit to detect the presence of a microbe or a virus.

44. The kit of claim 43, wherein the DNA internal segment and the two DNA arm segments are complementary to a contiguous universal microbial nucleic acid sequence.

45. The kit of claim 43, wherein at least one of the DNA arm segments is not hybridizable to the universal microbial nucleic acid sequence.

46. The kit of claim 43, wherein the DNA internal segment is about 7 to 19 nucleotides long; each DNA arm segment is about 5 to 9 nucleotides long; and each primer is about 10 to 20 nucleotides long.

47. The kit of claim 46, wherein the first DNA arm segments comprises at least one C or G nucleotide (s) at the 5' terminus and the second DNA arm segment comprises a similar number of complementary G or C nucleotide(s) at the 3' terminus, whereby the extension arms strongly hybridize by formation of at least one G:C pair.

48. The kit of claim 47, wherein the first DNA arm segment comprises 1 to 8 C or G nucleotide(s) at the 5' terminus and the second DNA arm segment comprises 1 to 8 complementary nucleotide(s) at the 3' terminus.

49. The kit of claim 48, wherein the first DNA arm segment further comprises at least one A or T nucleotide (s) vicinal to at least one C or G nucleotide(s) at the 5' terminus and the other arm comprises a similar number of complementary nucleotide(s) vicinal to at

least one G or C nucleotide (s) at the 3' terminus, whereby the hybridization of the extension arms is reinforced by formation of at least one A:T pair next to the at least one G:C pair.

50. The kit of claim 49, wherein the first DNA arm segment comprises 1 to 4 A or T nucleotide (s) vicinal to the C or G nucleotide at the 5' terminus and the second DNA arm segment comprises 1 to 4 complementary T or A nucleotide(s) vicinal to the G or C nucleotide(s) at the 3' terminus.

51. The kit of claim 43, wherein the first DNA arm segment has the nucleotide sequence 5'-GCTAG-3', and the second DNA arm segment has the nucleotide sequence 5'-CGATC-3'.

52. The kit of claim 44, wherein the DNA internal segment is hybridizably complementary to a conserved region of a ribosomal RNA (rRNA) operon encoding a 23S rRNA, and a first primer consists essentially of the nucleotide sequence 5'-GCTAAGGTCCCAAAGT-3', and a second primer consists essentially of the nucleotide sequence 5'-AGAACGCTCTCCTACC-3'.

53. The kit of claim 52, wherein the nucleic acid sequence of the DNA internal segment consists essentially of 5'-GCTTCTA-3'.

54. The kit of claim 52, wherein the nucleic acid sequence of the polynucleotide consists essentially of 5'-GGTGGCTGCTTCTAAGCCACC-3'.

55. The kit of claim 43, wherein the microbe is a bacterium, yeast, mold, or protist.

56. The kit of claim 43, wherein the DNA internal segment is complementary to a contiguous nucleic acid sequence which is specific to a class of microbes.

57. The kit of claim 56, wherein the class of microbes is bacteria, and the internal DNA segment comprises a contiguous bacterial-specific nucleic acid sequence.

58. The kit of claim 43, wherein the internal DNA segment is complementary to a contiguous universal or virus-specific nucleic acid sequence.

59. The kit of claim 43, wherein the polynucleotide further comprises a first and a second terminus, the first terminus being operatively connected to a fluorogen and the second terminus being operatively connected to a quencher.

60. The kit of claim 59, wherein the fluorogen is selected from the group consisting of pico green, fluorescein and edans.

61. The kit of claim 59, wherein the quencher is dabcy1.

62. The kit of claim 59, wherein the first terminus is the 5' terminus and the second terminus is the 3' terminus.

63. The kit of claim 59, wherein the nucleic acid sequence of the DNA internal segment consists essentially of 5'-GCTTCTA-3'.

64. The kit of claim 59, wherein the the nucleic acid sequence of the polynucleotide consists essentially of 5'-GGTGGCTGCTTCTAAGCCACC-3'.

65. The kit of claim 59, further comprising a first linker and a second linker operatively connecting the termini of the arm segments to the fluorogen and the quencher, respectively.

66. The kit in accordance with claim 65, wherein the linkers are comprised of C<sub>6</sub>-thio or C<sub>7</sub>-amino linkers.

67. The kit of claim 66, wherein each linker comprises about 1 to about 10 carbon atoms.

- 68. The kit of claim 43, wherein the primers and the polynucleotide are premixed.
- 69. The kit of claim 68, wherein the premix is in unitary amounts.
- 70. The kit of claim 68, wherein the premix is provided in bulk.
- 71. The kit of claim 43, further comprising a control selected from the group consisting of universal and specific microbial and viral nucleic acid sequences.
- 72. The kit of claim 43, further comprising PCR reagents.

## SEQUENCE LISTING

&lt;110&gt; Hunt-Wesson

<120> PCR TECHNIQUES FOR DETECTING MICROBIAL  
AND VIRAL CONTAMINANTS IN FOODSTUFFS

&lt;130&gt; FP02 42746

&lt;150&gt; 60/086,025

&lt;151&gt; 1998-05-18

&lt;160&gt; 137

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 38

&lt;212&gt; DNA

&lt;213&gt; Bacteria

&lt;400&gt; 1

cagcmgccc ggtaatwccc gtcaattcmt ttragttt

38

&lt;210&gt; 2

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Bacteria

&lt;400&gt; 2

gaagtcgtaa caaggcaagg catccaccgt

30

&lt;210&gt; 3

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Bacteria

&lt;400&gt; 3

gagtttgatc ctggcucact gctgcctccc gta

33

&lt;210&gt; 4

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Bacteria

&lt;400&gt; 4

agagtttgat cctggctcag aaggaggtga tccagccgca

40

&lt;210&gt; 5

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Bacteria

&lt;400&gt; 5

ccaaggggtc tgtggcgaca tttcaccggt aacaggattg

40

<210> 6  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 6  
tgctaatacc gtatgtgctt taaccgcgac cgggatgtca a 41

<210> 7  
<211> 35  
<212> DNA  
<213> Bacteria

<400> 7  
atggggatttc gtattaacga acttgaaccg atttg 35

<210> 8  
<211> 43  
<212> DNA  
<213> Bacteria

<400> 8  
accttggttac gacttcaccc cagagagttt gatcctggct cag 43

<210> 9  
<211> 31  
<212> DNA  
<213> Bacteria

<400> 9  
gatcctggct caggacgggc ggtgtgtaca a 31

<210> 10  
<211> 38  
<212> DNA  
<213> Bacteria

<400> 10  
aaactcaaat gaattgacgg gacgggcggt gtgtacaa 38

<210> 11  
<211> 42  
<212> DNA  
<213> Bacteria

<400> 11  
gatggaacca ccatttgwa gwacatcwat acawattcct gg 42

<210> 12  
<211> 45  
<212> DNA  
<213> Bacteria

<400> 12  
gatcctgtaa atggtgttgc aagtcccaat tattataact ttgat 45

<210> 13

<211> 46  
<212> DNA  
<213> Bacteria

<400> 13  
tgaaagctgg ctacaggaag gccggtttat gcagcaacgg acgtca 46

<210> 14  
<211> 37  
<212> DNA  
<213> Bacteria

<400> 14  
gcaactactg ttagttacct ccaagatatg ttttaac 37

<210> 15  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 15  
getgggcagc aaactgataa ctctccatca agctgtttgt tcgtccgccc 50

<210> 16  
<211> 46  
<212> DNA  
<213> Bacteria

<400> 16  
ttacggcggtt actatcctct ctaggtctcg gtcagatatg tgattc 46

<210> 17  
<211> 58  
<212> DNA  
<213> Bacteria

<400> 17  
cctctctata tgcacacgga gctcccagct atattgttga ctgcccggga cttcgacc 58

<210> 18  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 18  
tgtgtttgct tgtcttcagc atgcgcgggg tcatggaacg t 41

<210> 19  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 19  
ggcgacacat tataccgtgc ccgaattctg ttatatatgt c 41

<210> 20  
<211> 41

<212> DNA  
<213> Bacteria

<400> 20  
tctgtattgt ctttttcacc ttaatagcac ccggtacaag c 41

<210> 21  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 21  
ttttttctgt attttctttc ctttcaggca ggattacaac aattcacagc 50

<210> 22  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 22  
tcgccacacg ctgacgctga ccattacatg acctcgcttt agttcacaga 50

<210> 23  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 23  
gacggctgta ctgcagggtg tggcgatata ctttctgcag ggatgcaata 50

<210> 24  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 24  
ctccggagaa ctgggtgcat cttaccggag gagtaattac aaacctggca 50

<210> 25  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 25  
accctgtaaa cgaagtttgc gatctcatgc gactacttga c 41

<210> 26  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 26  
ttaatagcac ccggtacaag caggcctgac tcttcaaaag agaaaattac 50

<210> 27  
<211> 40  
<212> DNA



<213> Bacteria

<400> 27  
ccggtattac agaaatctga gtgcatgatg aatccagggt 40

<210> 28  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 28  
cagtctatta cagaactatg ccatactgat tgccgcaatt g 41

<210> 29  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 29  
gacaggattt gttaacagggt tccagttaca caatcaggcc 40

<210> 30  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 30  
acactggatg atctcagtgg ctgaatcccc ctccattatg 40

<210> 31  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 31  
ccatgacaac ggacagcagt tcctgtcaac tgagcacttt g 41

<210> 32  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 32  
aaatcgccat tcgttgacta cttcttgcca ttctggcaac tcgcatgca 50

<210> 33  
<211> 49  
<212> DNA  
<213> Bacteria

<400> 33  
cagtcgtcac tcactgggtt catcaggata ttctccccac tctgacacc 49

<210> 34  
<211> 41  
<212> DNA  
<213> Bacteria

<400> 34  
tttacgatag acttttcgac cacatataaa ttatttcgct c 41  
  
<210> 35  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 35  
aagaagatgt ttatggcggc cacgaatcag gttatgcctc 40  
  
<210> 36  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 36  
cattcacagt aaaagtggcc gggcgccctc cggtgagttc 40  
  
<210> 37  
<211> 48  
<212> DNA  
<213> Bacteria  
  
<400> 37  
ctgatcgaat ggctgccagg ctcccaacca gacgatagtt atcacgca 48  
  
<210> 38  
<211> 37  
<212> DNA  
<213> Bacteria  
  
<400> 38  
taatactcct gaacggcggt aggtgtcggc ttttctg 37  
  
<210> 39  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 39  
gaagagtccg tgggattacg agcgatgcag ctattaataa 40  
  
<210> 40  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 40  
tatggaattt cgccgatttt tgtttgctc cctgctgagg 40  
  
<210> 41  
<211> 44  
<212> DNA  
<213> Bacteria  
  
<400> 41

tgttacgtcc tgtagaaagc ccaaaactgc ctggcacagc aatt 44  
<210> 42  
<211> 38  
<212> DNA  
<213> Bacteria  
<400> 42  
ctggatggta tgggtgagggg aggccaacaa ttatttcc 38  
<210> 43  
<211> 41  
<212> DNA  
<213> Bacteria  
<400> 43  
gttccttgac cgcctttccg atacgccggt cagccaccct c 41  
<210> 44  
<211> 41  
<212> DNA  
<213> Bacteria  
<400> 44  
aagtgcgtca acgagcagct ttagtgcttt gtgaccatcg a 41  
<210> 45  
<211> 60  
<212> DNA  
<213> Bacteria  
<400> 45  
cataacgacg ccatcgcggc tctcaggaat ttgtgagcgc ttctgtcgtg gcagcgctaa 60  
<210> 46  
<211> 40  
<212> DNA  
<213> Bacteria  
<400> 46  
ctggagagac taacgggtgg attactgacg ctgattctgc 40  
<210> 47  
<211> 44  
<212> DNA  
<213> Bacteria  
<400> 47  
tggcaatcag cgtcaggtaa tggctaagag atcagcctat gtcc 44  
<210> 48  
<211> 36  
<212> DNA  
<213> Bacteria  
<400> 48  
gccaatggta aattagttct ccttaattgt ttttac 36

<210> 49  
<211> 30  
<212> DNA  
<213> Bacteria

<400> 49  
aggaggatga gatgaacttt attggctggc 30

<210> 50  
<211> 36  
<212> DNA  
<213> Bacteria

<400> 50  
cagcmgccgc ggtaatwcct ccataaaggt gaccct 36

<210> 51  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 51  
atgaatatga aaaaagcaac ttatacgcgga ccgaagccaa 40

<210> 52  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 52  
gctacagctg ggattgcggt ttatacgcgga ccgaagccaa 40

<210> 53  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 53  
actagcactc cagttgttaa ttatacgcgga ccgaagccaa 40

<210> 54  
<211> 37  
<212> DNA  
<213> Bacteria

<400> 54  
ttactgaggt agcragctta tacgcgaccg aagccaa 37

<210> 55  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 55  
cacgtgctac aatggataga gaatagtttt atgggattag 40

<210> 56

<211> 46  
<212> DNA  
<213> Bacteria

<400> 56  
catcgacggc aacctcggag aatacaatta cggttctcca ccattc 46

<210> 57  
<211> 34  
<212> DNA  
<213> Bacteria

<400> 57  
aacctatcca ggtgctccgc cacacttgag atat 34

<210> 58  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 58  
gcacatgcat tcaataaaga tgtcactgca tctccgtggt 40

<210> 59  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 59  
cggaggttcc gcaaaagatg cctccagagt gatcgatggt 40

<210> 60  
<211> 36  
<212> DNA  
<213> Bacteria

<400> 60  
cctaagacgc caatcgaaaa gcgcttgcaa ctgctc 36

<210> 61  
<211> 36  
<212> DNA  
<213> Bacteria

<400> 61  
cctaagacgc caatcgaaaa gcgcttgcaa ctgctc 36

<210> 62  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 62  
attgcgaaat ttggtacagc acttgagata tatgcaggag 40

<210> 63  
<211> 40

<212> DNA  
<213> Bacteria

<400> 63  
acaagctgca cctggtgcag tgacagcgtg tgtagtagca 40

<210> 64  
<211> 39  
<212> DNA  
<213> Bacteria

<400> 64  
caagcaacta cacctgcgcc gaaccttggt agcattcgt 39

<210> 65  
<211> 39  
<212> DNA  
<213> Bacteria

<400> 65  
acaagctgca cctggtgcag gasccttggt agcattcgt 39

<210> 66  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 66  
cgaatctaac ggctggcaca gcccaaatag tgtcaccgct 40

<210> 67  
<211> 42  
<212> DNA  
<213> Bacteria

<400> 67  
caaactgcta acacagctac tgcacttgaa ttgctgttat tg 42

<210> 68  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 68  
ccgggagctg ctaaaagcggg gccaaaccac cgaaaatacc 40

<210> 69  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 69  
gaagcacctt ttgacgaagc gctgggtgcta caggtgtttc 40

<210> 70  
<211> 40  
<212> DNA

<213> Bacteria

<400> 70  
aacaaggctt aactgtaaact actatagtca gctacaattg 40

<210> 71  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 71  
ttattaggat cgcgccaggc aaagaataac cgttggtcac 40

<210> 72  
<211> 39  
<212> DNA  
<213> Bacteria

<400> 72  
ttggaaacgg ttaaacgaag aaccttccca tcaaaaaca 39

<210> 73  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 73  
tcgcatcaaa ctgacaaacg gcaggtactc tataagtgcc 40

<210> 74  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 74  
gagagtcaac cagatcctaa accagatacc aaaagctatt ctcattttct 50

<210> 75  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 75  
atgaataaga gtcgatttat ttcatttacc cattctttgt tgtaagggtg 50

<210> 76  
<211> 50  
<212> DNA  
<213> Bacteria

<400> 76  
acacccaacg tatttagcaga gagccctgg tgcaggcatc atatcatacc 50

<210> 77  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 77  
gacataaaag ctaggaattt aaatcggatt aacattatcc 40  
  
<210> 78  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 78  
ctagttaggt aatatctcct taatgctata tcttataggg 40  
  
<210> 79  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 79  
tagataaagt taaaacaagc taacttacgg tggacccttc 40  
  
<210> 80  
<211> 61  
<212> DNA  
<213> Bacteria  
  
<400> 80  
ttcactattt gtaaaagtgt cagacccac ttactaatga atttttttat cgtaagccct 60  
t 61  
  
<210> 81  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 81  
atggcagcat cagcttgata tttccaataa ccaccggttt 40  
  
<210> 82  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 82  
ctagtgcatt tggtattcaa tgcattgaca ccatagtact 40  
  
<210> 83  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 83  
acggctatat acattcaatt tccatcgata atatacctaa 40  
  
<210> 84  
<211> 50  
<212> DNA  
<213> Bacteria



<400> 84  
agtatatagt gcaacttcaa ctaaaatcag cgttgtcttc gtcctcaata 50

<210> 85  
<211> 45  
<212> DNA  
<213> Bacteria

<400> 85  
gcgattgatg gtgatacggg tagccaagcc ttgacgaact aaagc 45

<210> 86  
<211> 48  
<212> DNA  
<213> Bacteria

<400> 86  
ctcagacggg atttgttagg cacgtctatc tctgtagccc ctattacg 48

<210> 87  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 87  
cgggcagatt ctagaccttc gcaccccaaa tagaactcga 40

<210> 88  
<211> 45  
<212> DNA  
<213> Bacteria

<400> 88  
cgggcagatt ctagacctcc tgcgatgac ttggagcatt cccac 45

<210> 89  
<211> 62  
<212> DNA  
<213> Bacteria

<400> 89  
tgaaataaag cagtcagggt gtgattctgc acacaaatca gggattctg cacacaaatc 60  
ag 62

<210> 90  
<211> 56  
<212> DNA  
<213> Bacteria

<400> 90  
ggaagcttcc aatgattcta agctggatgg gaattctcag gcgtccctac tggggt 56

<210> 91  
<211> 42  
<212> DNA  
<213> Bacteria

<400> 91  
ctgtcccttt tctgtcccc ggtctttagc tgcggcggtg gt 42

<210> 92  
<211> 48  
<212> DNA  
<213> Bacteria

<400> 92  
cgccgctcac tggggcactg gctgccagcc gttaagcgaa ccaccgc 48

<210> 93  
<211> 39  
<212> DNA  
<213> Bacteria

<400> 93  
ccgcggtaca ggttggcgca cgccaccac ttccgggcc 39

<210> 94  
<211> 38  
<212> DNA  
<213> Bacteria

<400> 94  
gaactgatg ataactgggg caattcaacc cacttcaa 38

<210> 95  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 95  
actcgatgat aactggggag cccccagtaa tccataaagg 40

<210> 96  
<211> 42  
<212> DNA  
<213> Bacteria

<400> 96  
ttaatgtgta cgctgcgagt gggagtattc atatgaagcg tc 42

<210> 97  
<211> 40  
<212> DNA  
<213> Bacteria

<400> 97  
ctattggta tgcgcaaagc tggaagtggg ttgaattgca 40

<210> 98  
<211> 42  
<212> DNA  
<213> Bacteria

<400> 98

aatgctgtct tcatttggag cgcaacatac atcacagcaa tc 42  
<210> 99  
<211> 38  
<212> DNA  
<213> Bacteria  
  
<400> 99  
aaagatattt ttgttcttgt gcagccagca cacgcggg 38  
<210> 100  
<211> 42  
<212> DNA  
<213> Bacteria  
  
<400> 100  
taagggtact atcgaggcgg acgtgaaatt aaccgtcaca ct 42  
<210> 101  
<211> 42  
<212> DNA  
<213> Bacteria  
  
<400> 101  
tcattggcaga acagcagtcga gactcatctt accattaaga ag 42  
<210> 102  
<211> 41  
<212> DNA  
<213> Bacteria  
  
<400> 102  
catggcagaa cagcagtcag actcatctta ccattaagaa g 41  
<210> 103  
<211> 36  
<212> DNA  
<213> Bacteria  
  
<400> 103  
attgagagcc tccgcgtggg tggcgaagcg gccctc 36  
<210> 104  
<211> 40  
<212> DNA  
<213> Bacteria  
  
<400> 104  
cttgttggtt tgaggccata tataaaaagtt ggcataaaca 40  
<210> 105  
<211> 42  
<212> DNA  
<213> Bacteria  
  
<400> 105  
caaattatga cagaatcctt cgagaaatat gacatggatt gc 42

<211> 34  
<212> DNA  
<213> Bacteria

<400> 113  
ctccagaatc atctccacag gtatacaaag tcag 34

<210> 114  
<211> 44  
<212> DNA  
<213> Bacteria

<400> 114  
cctccggccc ctgaatgcgg ctaattaaca gtggattcgt cggt 44

<210> 115  
<211> 7  
<212> DNA  
<213> Bacteria

<400> 115  
atcttcg 7

<210> 116  
<211> 7  
<212> DNA  
<213> Bacteria

<400> 116  
tagaagc 7

<210> 117  
<211> 21  
<212> DNA  
<213> Bacteria

<400> 117  
ggtggcttag aagcagccac c 21

<210> 118  
<211> 5  
<212> DNA  
<213> Bacteria

<400> 118  
gctag 5

<210> 119  
<211> 5  
<212> DNA  
<213> Bacteria

<400> 119  
cgatc 5

<210> 120  
<211> 16

<212> DNA  
<213> Bacteria

<400> 120  
gctaagggtcc caaagt 16

<210> 121  
<211> 16  
<212> DNA  
<213> Bacteria

<400> 121  
agaacgctct cctacc 16

<210> 122  
<211> 11  
<212> DNA  
<213> Bacteria

<400> 122  
cttagaagca g 11

<210> 123  
<211> 25  
<212> DNA  
<213> Bacteria

<400> 123  
cctgcacggg cgggtgtgtac gcagg 25

<210> 124  
<211> 28  
<212> DNA  
<213> Bacteria

<400> 124  
cccccgctcat gcggcttcgt tatggggg 28

<210> 125  
<211> 32  
<212> DNA  
<213> Bacteria

<400> 125  
gctgcctgcc cctgaactat ccaagaggca gc 32

<210> 126  
<211> 10  
<212> DNA  
<213> Bacteria

<400> 126  
agtcagccac 10

<210> 127  
<211> 20  
<212> DNA

<213> Bacteria

<400> 127  
atcgggctgt tatgatgata 20

<210> 128  
<211> 10  
<212> DNA  
<213> Bacteria

<400> 128  
aatcgggatg 10

<210> 129  
<211> 20  
<212> DNA  
<213> Bacteria

<400> 129  
atcactgggg gagctaatat 20

<210> 130  
<211> 10  
<212> DNA  
<213> Bacteria

<400> 130  
aggggtcttg 10

<210> 131  
<211> 23  
<212> DNA  
<213> Bacteria

<400> 131  
taagggtaaa gttgttgtga atc 23

<210> 132  
<211> 10  
<212> DNA  
<213> Bacteria

<400> 132  
agccagcgaa 10

<210> 133  
<211> 20  
<212> DNA  
<213> Bacteria

<400> 133  
catatccaga gcctctggat 20

<210> 134  
<211> 10  
<212> DNA  
<213> Bacteria

<400> 134	
gaaacgggtg	10
<210> 135	
<211> 23	
<212> DNA	
<213> Bacteria	
<400> 135	
gtagcctctt catcgctgc taa	23
<210> 136	
<211> 10	
<212> DNA	
<213> Bacteria	
<400> 136	
gtgacgtagg	10
<210> 137	
<211> 20	
<212> DNA	
<213> Bacteria	
<400> 137	
caccgccttt aacgccaaga	20

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/68</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/63112</b> <b>(43) International Publication Date:</b> 9 December 1999 (09.12.99)
<b>(21) International Application Number:</b> PCT/US99/10940 <b>(22) International Filing Date:</b> 18 May 1999 (18.05.99)  <b>(30) Priority Data:</b> 60/086,025 18 May 1998 (18.05.98) US Not furnished 17 May 1999 (17.05.99) US  <b>(71) Applicant (for all designated States except US):</b> HUNT-WESSON, INC. [US/US]; 1645 West Valencia Drive, Fullerton, CA 92833 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ROMICK, Thomas, L. [US/US]; 17637 Dandelion Lane, Chino Hills, CA 91709 (US). FRASER, Mark, S. [US/US]; 2331 E. Brookdale Place, Fullerton, CA 92831 (US).  <b>(74) Agents:</b> CRAFT, Jeffrey, F. et al.; Pretty, Schroeder & Poplawski, 19th floor, 444 South Flower Street, Los Angeles, CA 90071 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 16 March 2000 (16.03.00)
<b>(54) Title:</b> PCR TECHNIQUES FOR DETECTING MICROBIAL AND VIRAL CONTAMINANTS IN FOODSTUFFS  <b>(57) Abstract</b>  A method of detecting the presence of living or dead microorganisms and viruses in a sample comprises adding to a predetermined volume of a sample comprising nucleic acid-containing microbe(s) and/or virus(es), known amounts of a pair of primers binding to sequences upstream and downstream to a universal or specific microbial and/or viral nucleic acid sequence and polymerase chain reaction (PCR) reagents, cycling the mixture to amplify the universal or specific microbial and/or viral nucleic acid sequence; adding a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and a second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arms segments comprising nucleotide sequences such that they are hybridizably complementary to one another.		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

In.      nal Application No  
PCT/US 99/10940

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6    C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6    C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 13399 A (PUBLIC HEALTH RESEARCH INST OF) 18 May 1995 (1995-05-18) see esp. claims and figures ---	1-23, 43-72
X	EP 0 480 289 A (BOEHRINGER MANNHEIM GMBH) 15 April 1992 (1992-04-15) see examples ---	27, 33
X	TYAGI S ET AL: "MOLECULAR BEACONS: PROBES THAT FLUORESCCE UPON HYBRIDIZATION" NATURE BIOTECHNOLOGY, US, NATURE PUBLISHING, vol. 14, - March 1996 (1996-03) page 303-308 XP000196024 ISSN: 1087-0156 cited in the application the whole document ---	1-23, 43-72
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 December 1999

Date of mailing of the international search report

14/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Müller, F

## INTERNATIONAL SEARCH REPORT

Int. Application No  
PCT/US 99/10940

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 786 519 A (HUMAN GENOME SCIENCES INC) 30 July 1997 (1997-07-30) the whole document ---	
A	BARRY T ET AL: "THE 16S/23S RIBOSOMAL SPACER REGION AS A TARGET FOR DNA PROBES TO IDENTIFY EUBACTERIA" PCR METHODS & APPLICATIONS,US,COLD SPRING HARBOR LABORATORY PRESS, vol. 1, - 1991 page 51-56 XP000609831 ISSN: 1054-9803 the whole document ---	
A	EP 0 133 671 A (MILES LAB) 6 March 1985 (1985-03-06) see whole doc. esp. claims and page 1, 1.par. ---	
A	EP 0 229 943 A (MOLECULAR BIOSYSTEMS INC) 29 July 1987 (1987-07-29) the whole document ---	
A	WO 88 03957 A (GEN PROBE INC) 2 June 1988 (1988-06-02) page 161; claim 223 ---	
A	CHEN S ET AL: "A rapid, sensitive and automated method for detection of Salmonella species in food using AG-9600 AmpliSensor analyzer" JOURNAL OF APPLIED MICROBIOLOGY,GB,OXFORD, vol. 83, no. 3, - September 1997 (1997-09) page 314-321-321 XP002099375 ISSN: 1364-5072 the whole document ---	
A	WO 90 15157 A (GENE TRAK SYSTEMS) 13 December 1990 (1990-12-13) page 14 -page 16 ---	
X	WO 89 03891 A (CHIRON CORP) 5 May 1989 (1989-05-05) see esp. fig. 10-3 -----	36

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No  
PCT/US 99/10940

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9513399	A	18-05-1995	AU 697841 B	15-10-1998
			AU 1292195 A	29-05-1995
			CA 2176348 A	18-05-1995
			EP 0728218 A	28-08-1996
			JP 9504950 T	20-05-1997
			US 5925517 A	20-07-1999
EP 0480289	A	15-04-1992	DE 4038804 A	16-04-1992
			AT 150092 T	15-03-1997
			AT 122104 T	15-05-1995
			AU 655781 B	12-01-1995
			AU 8627591 A	28-04-1992
			CA 2052668 A	10-04-1992
			CA 2093647 A	10-04-1992
			DE 59105402 D	08-06-1995
			DE 59108610 D	17-04-1997
			DK 552203 T	02-10-1995
			WO 9206216 A	16-04-1992
			EP 0552203 A	28-07-1993
			ES 2101706 T	16-07-1997
			ES 2073771 T	16-08-1995
			FI 931586 A	07-04-1993
			IE 68525 B	26-06-1996
			IL 99663 A	31-12-1995
			JP 4258299 A	14-09-1992
			JP 6040839 B	01-06-1994
			JP 2673162 B	05-11-1997
			JP 6501157 T	10-02-1994
			NO 305995 B	30-08-1999
			NZ 240079 A	27-07-1993
			PT 99177 A,B	30-09-1992
			US 5753433 A	19-05-1998
EP 0786519	A	30-07-1997	CA 2194411 A	06-07-1997
			JP 9322781 A	16-12-1997
EP 0133671	A	06-03-1985	AU 3138784 A	07-02-1985
			CA 1231303 A	12-01-1988
			ES 534864 A	16-08-1985
			IL 72499 A	28-02-1989
			JP 60100056 A	03-06-1985
			US 5200313 A	06-04-1993
EP 0229943	A	29-07-1987	AT 66962 T	15-09-1991
			CA 1273552 A	04-09-1990
			DE 3681272 A	10-10-1991
			GR 3002735 T	25-01-1993
			JP 2033069 C	19-03-1996
			JP 7063400 B	12-07-1995
			JP 62157570 A	13-07-1987
			US 4996143 A	26-02-1991
WO 8803957	A	02-06-1988	AT 163680 T	15-03-1998
			AU 616646 B	07-11-1991
			AU 1041988 A	16-06-1988
			CA 1339871 A	19-05-1998
			DE 3752172 D	09-04-1998
			DE 3752172 T	02-07-1998

# INTERNATIONAL SEARCH REPORT

information on patent family members

In: International Application No  
PCT/US 99/10940

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 8803957 A		DK 413788 A	23-09-1988
		EP 0272009 A	22-06-1988
		ES 2112824 T	16-04-1998
		FI 883482 A	22-07-1988
		JP 10042880 A	17-02-1998
		JP 1503356 T	16-11-1989
		KR 9511719 B	09-10-1995
		NO 883248 A	23-09-1988
		PT 86204 A,B	01-12-1987
		US 5541308 A	30-07-1996
		US 5595874 A	21-01-1997
		US 5547842 A	20-08-1996
		US 5593841 A	14-01-1997
		US 5683876 A	04-11-1997
		US 5677127 A	14-10-1997
		US 5677128 A	14-10-1997
		US 5677129 A	14-10-1997
		US 5827651 A	27-10-1998
		US 5693468 A	02-12-1997
		US 5958679 A	28-09-1999
		US 5691149 A	25-11-1997
		US 5693469 A	02-12-1997
		US 5679520 A	21-10-1997
		US 5714321 A	03-02-1998
		US 5674684 A	07-10-1997
		US 5840488 A	24-11-1998
WO 9015157 A	13-12-1990	AT 127530 T	15-09-1995
		AU 5950690 A	07-01-1991
		CA 2031499 A	01-12-1990
		DE 69022180 D	12-10-1995
		DE 69022180 T	01-02-1996
		EP 0431149 A	12-06-1991
		JP 4500315 T	23-01-1992
		US 5401631 A	28-03-1995
WO 8903891 A	05-05-1989	AT 133713 T	15-02-1996
		DE 3854956 D	14-03-1996
		DK 294589 A	15-08-1989
		EP 0317077 A	14-05-1989
		ES 2083363 T	16-04-1996
		GR 3019046 T	31-05-1996
		IE 75211 B	27-08-1997
		JP 2749277 B	13-05-1998
		JP 7303498 A	21-11-1995
		JP 2109999 A	23-04-1990
		JP 2565552 B	18-12-1996
		US 5656731 A	12-08-1997
		US 5359100 A	25-10-1994
		US 5571670 A	05-11-1996
		US 5614362 A	25-03-1997
		US 5124246 A	23-06-1992
		US 5594118 A	14-01-1997
		US 5624802 A	29-04-1997

CORRECTED  
VERSION\*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12Q 1/68</b>		<b>A3</b>	(11) International Publication Number: <b>WO 99/63112</b> (43) International Publication Date: 9 December 1999 (09.12.99)
(21) International Application Number: PCT/US99/10940 (22) International Filing Date: 18 May 1999 (18.05.99) (30) Priority Data: 60/086,025 18 May 1998 (18.05.98) US Not furnished 17 May 1999 (17.05.99) US (71) Applicant (for all designated States except US): HUNT-WESSON, INC. [US/US]; 1645 West Valencia Drive, Fullerton, CA 92833 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROMICK, Thomas, L. [US/US]; 17637 Dandelion Lane, Chino Hills, CA 91709 (US). FRASER, Mark, S. [US/US]; 2331 E. Brookdale Place, Fullerton, CA 92831 (US). (74) Agents: CRAFT, Jeffrey, F. et al.; Pretty, Schroeder & Poplawski, 19th floor, 444 South Flower Street, Los Angeles, CA 90071 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. (88) Date of publication of the international search report: 16 March 2000 (16.03.00)	
(54) Title: PCR TECHNIQUES FOR DETECTING MICROBIAL AND VIRAL CONTAMINANTS IN FOODSTUFFS			
(57) Abstract <p>A method of detecting the presence of living or dead microorganisms and viruses in a sample comprises adding to a predetermined volume of a sample comprising nucleic acid-containing microbe(s) and/or virus(es), known amounts of a pair of primers binding to sequences upstream and downstream to a universal or specific microbial and/or viral nucleic acid sequence and polymerase chain reaction (PCR) reagents, cycling the mixture to amplify the universal or specific microbial and/or viral nucleic acid sequence; adding a polynucleotide comprising a DNA internal segment that is hybridizably complementary to at least a portion of the universal or specific nucleic acid sequence; and a first and second DNA arm segment adjoining the DNA internal segment, the first DNA arm segment ending in a 5' terminus and the second DNA arm segment ending in a 3' terminus, the arm segments comprising nucleotide sequences such that they are hybridizably complementary to one another. Also disclosed are polynucleotides and kits useful for practicing the method.</p>			

\*(Referred to in PCT Gazette No. 17/2000, Section II)

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						